

# Reactive Scattering between Metastable Helium and Magneto-Optically Trapped Lithium Atoms

J. Grzesiak, S. Hofsäss, F. Stienkemeier, M. Mudrich, K. Dulitz<sup>1</sup>

Department of Physics, University of Freiburg, Hermann-Herder-Str. 3, 79104 Freiburg, Germany

The experimental study of Penning ionization reactions, i.e., the reactive scattering of metastable rare gas atoms with neutral species, has recently attracted a lot of attention through the observation of orbiting resonances at low collision energies [1].

Our project is aimed at a detailed study of reactive Penning collisions between quantum-state-selected lithium atoms (Li) and metastable helium atoms (He\*) in order to elucidate the influence of spin polarization on the reaction rate. Our setup consists of a cryogenically cooled source for the production of velocity-tunable, supersonic beams of metastable helium atoms and a magneto-optical trap (MOT) for ul-

tracold Li atoms which serves as a stationary scattering target. By selectively switching off the MOT laser beams, we can selectively populate one of the two hyperfine components of the  $^2S_{1/2}$  electronic ground state of Li. In this contribution, we will present first experimental results for reactive Penning collisions between He\* and Li at different collision energies, including a detailed discussion of the supersonic beam and MOT characteristics.

## References

- [1] A. B. Henson, S. Gersten, Y. Shagam, J. Narevicius, E. Narevicius 2012 *Science* **338** 234

---

<sup>1</sup>E-mail: [katrin.dulitz@physik.uni-freiburg.de](mailto:katrin.dulitz@physik.uni-freiburg.de)