



International Mountain Conference 2019, Innsbruck

A multi-year field experiment for studying the importance of different abiotic and biotic drivers of tree seedling recruitment in an alpine treeline ecotone

Esther R. Frei^{1,2}, Peter Bebi²,
Christian Rixen², Carissa Brown³

¹Swiss Federal Institute for Forest, Snow and Landscape Research WSL

²WSL Institute for Snow and Avalanche Research SLF


³Department of Geography, Memorial University, St. John's, Canada



Global Treeline Range Expansion Experiment Initiative (G-TREE)

- Globally distributed multi-year field experiments to test multiple drivers of seedling recruitment at treelines
- Common experimental protocol
- Research aims:
 - Disentangling seed vs. substrate limitations
 - Predictions for future treeline expansion


16 sites
worldwide



<http://treelineresearch.com/>

Swiss G-TREE experiment

Experimental sites near Davos,
Switzerland (established in 2013)



Study species



Picea abies




Larix decidua

- low and high elevation provenances of both species

Treatments

- seeding (2013 and 2014)
- soil scarification
- herbivore exclosures


Seedling emergence



- No natural recruitment at and above treeline
- Emergence increased with elevation but varied with seeding year
- Dwarf shrubs and dense understorey vegetation reduced emergence

Frei *et al.* (2018) *SciRep*



Seedling survival



- Winter survival was higher at treeline than at the alpine site but with larger differences between sites in 2014
- Higher survival on scarified plots
- 1st winter survival is a bottleneck for recruitment


Frei *et al.* (2018) *SciRep*

Seedling survival



- Winter survival was higher at treeline than at the alpine site but with larger differences between sites in 2014
- Higher survival on scarified plots
- 1st winter survival is a bottleneck for recruitment

Frei *et al.* (2018) *SciRep*



Synthesis

- Tree seedlings can emerge and establish at and well above the current treeline when viable seeds reach suitable microsites for germination
- Multiple abiotic and biotic drivers shape early seedling recruitment in the treeline ecotone
- Global results support findings of the Swiss study: strong seed and substrate limitations at treelines worldwide

Acknowledgements

G-TREE partners, Eva Bianchi, Melissa Dawes, Martin Heggli, Alessio Pataglia, Ueli Schmid, Moritz Wanner, Lea Voegelé, Sonja Wipf, Giulietta Bernareggi, Werkstatt SLF, Alpgenossenschaft Davos, field crews 2013-2018