

Characteristics of alpine plants and soils along an elevational gradient, Northern Selkirk Mountains, British Columbia

Author: Cole Brachmann¹


Supervisors: David S Hik¹, Guillermo Hernandez Ramirez²

¹Department of Biological Sciences, University of Alberta

²Department of Renewable Resources, University of Alberta

Survival and growth of alpine plant species in recently-deglaciated terrain

- Current alpine plant community and soil property distributions
- Transplantation of alpine species into recently-deglaciated terrain
- Potential for alpine species to shift upslope as climate change continues



Controlling factors of soil CH₄ uptake in temperate alpine

- Soil-atmosphere methane fluxes understudied in Canadian alpine
- Methane uptake related to total carbon, $\delta^{15}\text{N}$ and moisture
- Average uptake of:
 $-112 \pm 12 \mu\text{g CH}_4\text{-C m}^{-2} \text{ hr}^{-1}$



Controlling factors of soil CH₄ uptake in temperate alpine

- Soil-atmosphere methane fluxes understudied in Canadian alpine
- Methane uptake related to total carbon, $\delta^{15}\text{N}$ and moisture
- Average uptake of:
 $-112 \pm 12 \mu\text{g CH}_4\text{-C m}^{-2} \text{ hr}^{-1}$

