

Biosynthesis and Structure of Aeruginoside 126A and 126B, Cyanobacterial Peptide Glycosides Bearing a 2-Carboxy-6-Hydroxyoctahydroindole Moiety

Keishi Ishida,^{1,6,8} Guntram Christiansen,^{1,2,8} Wesley Y. Yoshida,² Rainer Kurmayer,³ Martin Welker,⁴ Nativitat Valls,⁵ Josep Bonjoch,⁵ Christian Hertweck,⁶ Thomas Börner,¹ Thomas Hemscheidt,^{2,7,*} and Elke Dittmann^{1,*}

¹Institute of Biology, Humboldt University Berlin, 10115 Berlin, Germany

²Department of Chemistry, University of Hawaii at Manoa, Honolulu, HI 96822, USA

³Institute for Limnology, Austrian Academy of Science, 5310 Mondsee, Austria

⁴Institute of Chemistry, Technical University Berlin, 10587 Berlin, Germany

⁵Faculty of Pharmacy, University of Barcelona, 08028 Barcelona, Spain

⁶Department of Biomolecular Chemistry, Leibniz-Institute for Natural Product Research and Infection Biology (HKI), 07745 Jena, Germany

⁷Cancer Research Center of Hawaii, Honolulu, HI 96813, USA

⁸These authors contributed equally to this work.

*Correspondence: hemschei@hawaii.edu (T.H.), elke.dittmann@rz.hu-berlin.de (E.D.)

DOI 10.1016/j.chembiol.2007.04.006

SUMMARY

Aeruginosins represent a group of peptide metabolites isolated from various cyanobacterial genera and from marine sponges that potently inhibit different types of serine proteases. Members of this family are characterized by the presence of a 2-carboxy-6-hydroxyoctahydroindole (Choi) moiety. We have identified and fully sequenced a NRPS gene cluster in the genome of the cyanobacterium *Planktothrix agardhii* CYA126/8. Insertional mutagenesis of a NRPS component led to the discovery and structural elucidation of two glycopeptides that were designated aeruginoside 126A and aeruginoside 126B. One variant of the aglycone contains a 1-amino-2-(*N*-amidino- Δ^3 -pyrrolinyl)ethyl moiety at the C terminus, the other bears an agmatine residue. In silico analyses of the aeruginoside biosynthetic genes *aerA*–*aerI* as well as additional mutagenesis and feeding studies allowed the prediction of enzymatic steps leading to the formation of aeruginosides and the unusual Choi moiety.

INTRODUCTION

Cyanobacteria provide a broad range of biologically active compounds, many of which are highly intriguing from a structural point of view [1]. In particular, cyanobacterial peptides are often composed of highly unusual amino acids. A good example is the group of aeruginosins, which are characterized by an *N*-terminal derivative of an aryl lactic acid, a 2-carboxy-6-hydroxyoctahydroindole (Choi) moiety, and an arginine derivative at the C-terminal posi-

tion [2]. Different variants of aeruginosin have been isolated from the planktonic cyanobacterial genera *Microcystis*, *Oscillatoria* (*Planktothrix*), *Nodularia*, and from marine sponges [2–11]. Aeruginosins are potent inhibitors of serine proteases with different specificities for trypsin and thrombin, respectively. Two specific aeruginosins have also been crystallized bound to their protease targets trypsin and thrombin, respectively, in order to assess the structure-activity relationships [12, 13].

The majority of cyanobacterial peptides is synthesized by modular biosynthetic assembly lines, so-called nonribosomal peptide synthetases (NRPS) [1, 14]. A NRPS module is minimally composed of an amino acid-activating adenylation (A) domain, a peptidyl carrier protein (PCP) domain, and a condensation (C) domain [15]. In the last decade, a number of such NRPS systems have been described from cyanobacteria. Biosynthesis genes for the production of the widespread cyanobacterial hepatotoxin microcystin were detected and characterized in *Microcystis*, *Planktothrix*, and *Anabaena*, three distant cyanobacterial genera that form blooms in freshwater habitats [16–19]. The microcystin pathway exemplifies the prevalence of mixed NRPS/polyketide synthase (PKS) pathways, which is a major theme of cyanobacterial biosyntheses. This biosynthetic principle is also seen in several pathways that have been reported from the marine cyanobacterium *Lyngbya majuscula*, namely the barbamide, jamaiamide, and curacin pathways [20–22]. Natural product biosyntheses from cyanobacteria have further attracted attention because of their unique tailoring enzymes, including diverse forms of halogenases and aromatic prenyl transferases [20, 23, 24]. These fascinating properties of cyanobacterial NRPS/PKS pathways make them particularly interesting for pathway engineering approaches. It was therefore tempting to investigate the natural biosynthetic machinery of the structurally unique aeruginosin and its core moiety Choi.




Figure 1. Inactivation of NRPS Presumably Involved in Aeruginosin Biosynthesis by Homologous Recombination and HPLC Analysis of Wild-Type and Mutant

(A) Schematic representation of Δ NRPS-D-Leu knockout plasmids for the insertional mutagenesis of *P. agardhii* CYA126/8. The chloramphenicol resistance gene cassette (Cmr) is highlighted in black.

(B) PCR amplification with the DNA from wild-type (WT) and Δ NRPS-D-Leu mutant (MT) with primers amplifying the region flanking the insertion site.

(C) HPLC-DAD profile of extract from *Planktothrix agardhii* CYA126/8 wild-type.

(D) HPLC-DAD profile of extract from Δ NRPS-D-Leu mutant. The peaks at min 9.07 (1) and 10.56 (2) represent putative aeruginosins, the peaks at min 26.6 (3) and 33.8 (4) [p-Asp]-microcystin-RR and [p-Asp]-microcystin-LR, respectively.

Here, we report the cloning, sequencing, and mutational analysis of a gene cluster encoding aeruginosin biosynthesis in *Planktothrix agardhii* CYA126/8 as well as the full structural elucidation of two previously unreported glycosylated aeruginosin variants, aeruginosides 126A and 126B.

RESULTS AND DISCUSSION

Identification of the Aeruginoside Gene Cluster via Gene Disruption

A candidate gene cluster for the biosynthesis of the aeruginosides was identified in the course of a search for NRPS genes by a degenerate PCR approach in *Planktothrix* CYA126/8 [16]. Subsequently, individual PCR fragments were cloned and sequenced and used to screen a phage-mide library of the strain. As a result, a gene fragment was identified encoding a NRPS module comprising an A domain, a PCP domain, and an epimerization (E) domain (Figure 1A). Analysis of the specificity conferring residues of the amino acid binding pocket [25] and comparison with characterized sequences in the database revealed a putative specificity of the A domain for leucine (Figure 1A and Table S1 available with this article online). Furthermore, the presence of an epimerization domain suggested that the module could be responsible for the incorporation of a D-leucine moiety into a cyanopeptide structure. Leucine is common in several cyanobacterial peptide families; however, the D stereoisomer is characteristic for some aeruginosin variants that have been described from *Microcystis* [6]. Although *Planktothrix* CYA126 was known to produce a number of different peptides including putative aeruginosin variants [16], no detailed structural information on aeruginosins from CYA126 was available. In order to confirm the proposed involvement of the gene fragment in aeruginosin biosynthesis, an insertional mutagenesis of the gene in *Planktothrix* CYA126 was initiated by using a homologous recombination strategy. For this purpose a chloramphenicol resistance cassette was

inserted into the NRPS gene fragment, and the resulting construct was introduced into the *Planktothrix* cells by electroporation. Following selection in liquid medium, mutant clones were obtained. Subsequent PCR using primers distal to the insertion region revealed the successful insertion of the resistance cassette and the homozygosity of the mutants (Figure 1B).

Identification and Structural Elucidation of Aeruginosides 126A and 126B

The composition of the peptide fraction of wild-type cells and of mutant cells cultured in the presence of chloramphenicol was compared by HPLC-DAD analysis. This analysis revealed that in the extract from the mutant, three peaks were missing (Figures 1C and 1D and data not shown). MALDI-TOFMS analysis for two of these indicated molecular masses of 714 and 690, respectively (data not shown). These masses did not match those of any known aeruginosin-type molecules, which suggested that these compounds might be new. Thus, the peaks missing in the mutant were collected from large-scale laboratory cultures of the wild-type. Two metabolites, aeruginoside 126A (1) and aeruginoside 126B (2), were obtained in sufficient quantity, 0.8 mg and 0.4 mg, respectively, to allow for a full spectroscopic characterization by NMR.

Positive ion HRESI-TOFMS analysis of 1 yielded a pseudomolecular ion (M+H)⁺ at *m/z* 715.4056, suggesting a molecular formula of C₃₆H₅₅N₆O₉ (calc. 715.4031, Δ = 2.5 mmu). Inspection of the ¹H and ¹³C NMR spectra recorded in DMSO-*d*₆ indicated the presence of two conformers in the ratio of 3:1. Only the data for the major conformer are displayed in Table 1 and will be discussed in the following. The ¹³C NMR spectrum showed 34 signals two of which were assigned to methyl groups on the basis of the analysis of gHSQC data, 11 to methylenes, 15 to methines, and 6 to quaternary carbon atoms. This suggested that the molecule contained a symmetrical substructure, which was readily assigned to a phenyl group on the basis of the ¹H NMR spectrum. This aromatic ring accounted

only for 4 of the 13 degrees of unsaturation. From the ^{13}C NMR data, five of the remaining double-bond equivalents were identified in the form of three carbonyl resonances, a guanidine carbon atom, plus one C-C double bond. Thus, **1** had to contain four additional rings (Figure S1).

The identification of individual residues was accomplished by means of 1D TOCSY spectra. In this fashion, a leucine residue and a β -substituted lactic acid were readily identified. The large geminal coupling constant for the protons on C-3 of the latter suggested the presence of a π -system in the β -position. Eventually, this residue was identified as a phenyl lactic acid (Plac) on the basis of gHMBC data and chemical shift considerations. Thus, the assignment of the latter as an α -hydroxy acid rather than an amino acid is based on the chemical shift for C-2 (72.02 ppm) and H-2 (4.11 ppm). Further inspection of the ^{13}C NMR spectrum of **1** provided evidence for a 2-carboxy-6 α -hydroxy-octahydroindole (Choi) subunit with excellent agreement of the chemical shifts observed for **1** and those reported in the literature. The only significant chemical-shift difference noted was for the hydroxymethine carbon atom of **1** at $\delta = 70.05$ ppm, which was shifted downfield ($\Delta\delta \approx 6$ ppm) in comparison to Choi residues in other aeruginosin-type molecules ($\delta = 63.8$ to 64.1 ppm) [4, 6, 8]. Since the correct assignment of the relative stereochemistry of the Choi residue has proven problematic in the past, we paid particular attention to this aspect. The *cis* stereochemistry of the ring junction follows from the coupling constant $^3J = 6.2$ Hz observed for H-7a. Confirmation of this stereochemistry by NOE was not possible because of the overlap of the resonances for H-7 $_{eq}$ and H-3a. All expected NOE's were observed (see Figure S2). Finally, the arginine mimetic characteristic of the aeruginosin class of compounds was identified as a 1-amino-2-(*N*-amidino- Δ^3 -pyrrolinyl)-ethyl (Aeap) residue on the basis of gCOSY and gHMBC data using the NH and the olefinic methine proton resonances as starting points. This assignment was corroborated through chemical-shift comparison with literature values [26, 27], which were in excellent agreement.

The subunits of **1** identified thus far account for all but five carbon atoms, five oxygen atoms, and ten protons. This elemental composition of the remainder suggested the presence of a carbohydrate moiety in **1**, and a proton resonance at 4.81 ppm was tentatively assigned to the anomeric proton of the putative sugar. In the HSQC spectrum, the signal for this proton correlates to a resonance at 96.5 ppm, a typical value observed for acetal carbon atoms. Further analysis of gHMBC and gHSQC data led to the identification of three additional hydroxymethine and one hydroxymethylene carbon atom. The relative stereochemistry of the sugar was established on the basis of 1D TOCSY data. The large 3J coupling constants ($^3J_{2,3} = 9.4$ Hz; $^3J_{3,4} = 8.8$ Hz; $^3J_{4,5ax} = 10.6$ Hz) observed for the hydroxymethine protons on C-2, C-3, and C-4 of the sugar suggested an all-axial arrangement of these protons in a six-membered ring. Thus, the sugar residue in **1** was identified as a xylopyranoside. The small coupling constant between the anomeric proton and H-2 ($^3J = 3.4$ Hz) was

indicative of H-1 being equatorial and hence of the presence of an α -glycosidic linkage. The observation of a gHMBC correlation from H-1 of the sugar to C-6 ($\delta = 70.05$ ppm) of the Choi residue indicates the site of attachment of the sugar to the aglycone. This connectivity also explains the downfield shift of the signal for this carbon atom, as noted earlier, relative to that found in Choi units bearing a free hydroxyl.

The sequence of the amino acids in the peptide was determined by means of gHMBC correlations from the exchangeable NH proton of the leucine unit to C-1 of Plac and from the NH-1 resonance of Aeap to the carboxyl resonance of Choi. No HMBC correlations were observed in support of the bond between C-1 of leucine and the nitrogen atom of Choi. However, unambiguous evidence for a connection between these units is furnished by very strong NOE correlations between the aminomethine proton of leucine (4.50 ppm) and H-7a as well as H-7 $_{eq}$ of the Choi unit.

Positive ion HRESI-TOFMS analysis of **2** yielded a pseudomolecular ion ($\text{M}+\text{H}$) $^+$ at 691.4051, suggesting a molecular formula of $\text{C}_{34}\text{H}_{54}\text{N}_6\text{O}_9$ ($\Delta = 2.0$ mmu). As in the case of **1**, analysis of the NMR spectra of **2** revealed the presence of two conformers. Chemical-shift comparison of the NMR spectra of **1** and **2** revealed excellent agreement ($\Delta\delta$ (^{13}C) ≤ 0.1 ppm, $\Delta\delta$ (^1H) ≤ 0.05 ppm) for all but the arginine mimetic portion of the peptides. Analysis of gHMBC and 1D TOCSY data then established that **2** was bearing an agmatine rather than an Aeap moiety. Thus, the structure of **2** closely resembles the recently revised structure of aeruginosin 205 [28]. The absolute stereochemistry of the phenyl lactic acid and the leucine moiety of **1** was determined after acid hydrolysis by chromatography on chiral stationary phase and comparison of retention times to those of authentic standards. The retention times of Plac from **1** matched that of *D*-Plac and that of leucine matched the retention time of the *D*-leucine standard. The absolute configuration of the xylose residue could not be established (Table 1 and Figure 2). Aeruginoside 126A weakly inhibited porcine pancreas trypsin and bovine plasma thrombin with IC_{50} values of 67 and 30 $\mu\text{g}/\text{ml}$, respectively, compared to aeruginosin 98-A [5] as a positive control (0.07 and 2.5 $\mu\text{g}/\text{ml}$). Aeruginoside 126A did not inhibit human plasma plasmin at a concentration of 100 $\mu\text{g}/\text{ml}$, compared to aeruginosin 98-A [5] as a positive control with IC_{50} value of 2.7 $\mu\text{g}/\text{ml}$.

Analysis of the Organization of the Aeruginoside Gene Cluster and Flanking Regions

In order to obtain sequence information about the entire gene cluster encoding aeruginosin biosynthesis, the original fragment of the aeruginoside gene cluster was used as a probe to screen a genomic phagemid library of *PlanktOTHrix* CYA126/8. Subsequently 35 kb were sequenced from overlapping phagemid clones spanning the aeruginoside biosynthesis gene cluster and flanking regions. Sequence analysis of the aeruginosin (*aer*) region revealed the presence of 19 ORFs, nine of which could be assigned to aeruginoside biosynthesis and therefore designated

Table 1. NMR Spectral Data for Aeruginoside A (1) in DMSO- d_6

Unit	C/H No.	δ_H (J in Hz)	δ_C	HMBC
Plac	1		173.2	Plac H-2, H-3, Leu NH
	2	4.11, dd (8.8, 3.9)	72.02	Plac H-3 α , H-3 β
	3 α	2.95, dd (13.7, 3.9)	40.4	Plac H-2, H-5/9
	3 β	2.65, dd (13.7, 8.8)		
	4		138.4	Plac H-2, H-3 α , H-3 β , H-6/8
	5/9	7.21, m	129.3	Plac H-3 α , H-3 β , H-7, H-9/5
	6/8	7.24, m	127.9	Plac H-8/6
	7	7.16, m	125.9	Plac H-5/9
	Leu	1		169.9
2		4.50, ddd (9.0, 7.9, 4.8)	48.6	Leu H-3 α
3 α		1.46, m	41.0	Leu H-4, H-5, H-5'
3 β		1.34, m		
4		1.48, m	24.1	Leu H-3 α , H-5, H-5'
5		0.86, d (6.5)	21.8	Leu H-3 α , H-3 β , H-5'
5'		0.84, d (6.4)	23.3	Leu H-3 β , H-5
NH		7.76, d (7.9)		
Choi	1		171.4	Choi H-2, Aeap NH, H-1
	2	4.18, dd (9.0, 7.8)	59.8	Choi H-7a
	3 α	2.04, ddd (13.0, 7.8, 6.8)	30.5	Choi H-2
	3 β	1.78, ddd (13.0, 13.0, 9.0)		
	3a	2.28, m	36.1	
	4 _{ax}	2.08, m	19.5	
	4 _{eq}	1.47, m		
	5 _{eq}	1.56, m	24.5	
	5 _{ax}	1.49, m		
	6	3.84, m	70.05	Xyl H-1
	7 _{eq}	2.27, brdd (14.0, 4.8)	29.2	
	7 _{ax}	1.57, brdd (14.0, 11.6)		
	7a	4.15, ddd (11.6, 6.2, 4.8)	54.11	Choi H-3 α , H-4 _{eq} , H-7 _{eq}
Aeap	1 α	3.21, m	36.6	Aeap NH, H-2 α , H-2 β
	1 β	3.17, m		
	2 α	2.28, m	28.2	Aeap H-1, H-4
	2 β	2.26, m		
	3		136.2	Aeap H-1, H-2 α , H-2 β , H-4, H-7
	4	5.61, brs	119.1	Aeap H-2 α , H-2 β , H-5
	5	4.10, m	54.08	
	6		154.2	
7	4.09, m	55.3	Aeap H-4	
NH	7.74, t (5.8)			
Xyl	1	4.81, d (3.4)	96.5	Xyl H-3, H-5 _{ax}
	2	3.21, dd (9.4, 3.4)	71.94	Xyl H-3
	3	3.40, dd (9.4, 8.8)	73.4	Xyl H-1, H-2, H-5 _{ax} , H-5 _{eq}

Table 1. Continued

Unit	C/H No.	δ_H (J in Hz)	δ_C	HMBC
4		3.28, m	70.03	Xyl H-3, H-5 _{ax}
5 _{eq}		3.39, m	62.1	Xyl H-1
5 _{ax}		3.34, dd (10.6, 10.6)		

aerA-aerI. The function of ten additional ORFs remained obscure (Table 2 and Figure 3A). Nine of the Aer proteins are encoded in the *aerA-I* cluster. Upstream of the *aer* region, a gene was transcribed in the opposite direction encoding a protein with 51% similarity to the cyanobacterial circadian clock protein KaiA that was not considered to be involved in aeruginoside production (ORF1) (Figure 3A). Further upstream, a second ORF encoded a protein with no significant similarity to characterized proteins and was also not assigned to the biosynthesis (ORF2) (Figure 3A). A third ORF encoded between the *aerB* and *aerC* genes did not show any similarity to biosynthetic proteins and was thus not expected to be involved in aeruginoside biosynthesis (ORF3) (Figure 3A). Downstream of the *aerA-I* gene cluster, seven additional genes were sequenced. An overview of the deduced functions of these (ORF4–ORF10) is given in Table 2. One of these genes encodes a protein with 46% similarity to a glycosyltransferase and was designated *aerI*. This tailoring activity is needed for the modification of the aeruginosin base structure and was therefore attributed to aeruginoside biosynthesis. The sequence of the *aer* gene cluster and flanking regions has been deposited in the EMBL database (accession number AM071396). In the following section, the individual Aer proteins are analyzed in detail. A schematic representation of the predicted aeruginoside biosynthesis pathway is shown in Figure 3B.

In Silico Analysis of AerA–J Proteins

The first protein encoded by the *aer* gene cluster, AerA, represents an unusual PKS-like module comprising a NRPS A-domain, a ketoreductase (KR) domain, and an acyl carrier protein (ACP) domain. Analysis of the substrate binding pocket showed closest similarity to the A domain of the microcystin biosynthesis protein McyG that is known to activate diverse phenylpropanoid starter units in vitro [29] (Table S1). It is therefore proposed that AerA activates phenylpyruvate, which is reduced by the

ketoreductase domain to yield the Plac moiety of aeruginoside. The rare phenylpyruvate starter [30] is likely provided in sufficient amounts by the shikimate pathway.

Further downstream, the *aerB* gene encodes a NRPS module comprising a C, A, PCP, and E domain. This gene was initially inactivated resulting in the loss of aeruginoside production. As mentioned above, the substrate activating binding pocket shows close similarity to leucine activating domains (Table S1). The AerB module is thus expected to incorporate D-leucine into the peptide structure. AerC contains signature sequences of ring-hydroxylating dioxygenases and a conserved Rieske [2Fe-2S] domain (Table 2).

AerD, E, and F show 58%, 45%, and 55% amino acid sequence similarity to BacA, BacB, and BacC of bacilysin biosynthesis. These three enzymes are involved in the biosynthesis of the anticapsin moiety [31] of bacilysin (L-alanyl-(2,3-epoxycyclohexan-4-one)L-alanine). An alignment of the sequences of the Bac enzymes from *Bacillus amyloliquefaciens* and *Bacillus subtilis* [31] and of the corresponding Ywf enzymes encoded in the completely sequenced genome of *Bacillus subtilis* subsp. *subtilis* str. 168 [32] with AerD, AerE, and AerF, respectively, is provided as Supplemental Data (Figure S3). Feeding studies and studies of knock out mutants of chorismate mutase have revealed prephenate as precursor of the anticapsin moiety [33]. A motif scan of AerD using protein family analysis software (<http://www.sanger.ac.uk/Software/Pfam/search.shtml>) showed a conserved prephenate dehydratase signature. Prephenate is a known intermediate of phenylalanine and of tyrosine biosynthesis in many organisms. We conclude that AerD is involved in the biosynthesis of Choi in vivo. BacB and AerE, respectively, contain a conserved cupin domain. Although the role of this domain is not well understood, it is conserved among several dioxygenases [34]. BacB could perform the epoxidation step in anticapsin synthesis. It is not clear, what function could be fulfilled by the related enzyme AerE in Choi biosynthesis.




Figure 2. Structures of Aeruginoside 126A (1) and 126B (2)

D-Plac, D-phenyl lactic acid; Choi, 2-carboxy-6-hydroxyoctahydroindole; Aeap, 1-amino-2-(N-amidino- Δ^3 -pyrrolyl)ethyl; Agm, agmatin; Xyl, xylose.

Table 2. Deduced Functions of ORFs in the Aeruginoside Biosynthetic Gene Cluster

Protein	Amino Acids	Deduced Function	Sequence Similarity	Identity/Similarity (AA Length)	Accession Number
AerA	1416	PKS (A KR ACP)	BarE (<i>Lyngbya majuscula</i>)	44%/64% (587)	AAN32979
AerB	1598	NRPS (C A T E)	NosC (<i>Nostoc</i> sp. GSV224)	54%/71% (944)	AAF17280
AerC	736	Oxygenase	Putative diaminopimelate decarboxylase (<i>Burkholderia mallei</i> 10229)	30%/49% (707)	ZP00431845
AerD	202	Decarboxylase	BacA (<i>Bacillus amyloliquefaciens</i> AAM90573)	39%/58% (191)	AAM90573
AerE	213	Unknown	BacB (<i>Bacillus subtilis</i>)	26%/45% (208)	AAM90569
AerF	264	Reductase	BacC (<i>Bacillus subtilis</i>)	36%/55% (256)	AA65204
AerG	1622	NRPS (C A T C T)	McyB (<i>Microcystis aeruginosa</i>)	47%/64% (927)	AAE09608
AerH	308	Dioxygenase	Thymine dioxygenase (<i>Rhodotorula glutinis</i>)	30%/50% (517)	AAU12179
AerI	418	Glycosyltransferase	Spore coat protein SA (<i>Bacillus subtilis</i>)	26%/46% (336)	P46915
ORF1	248	Circadian clock protein	KaiA (<i>Synechococcus elongatus</i>)	31%/51% (227)	1R8JA
ORF2	126	Unknown	no similarity		
ORF3	288	Unknown	no similarity		
ORF4	229	Unknown	no similarity		
ORF5	170	Unknown	Putative reverse transcriptase (<i>Nostoc</i> sp. PCC7120)	70%/83% (157)	BAB77479
ORF6	428	Unknown	Retron-type reverse transcriptase (<i>Anabaena variabilis</i> ATCC29413)	58%/74% (415)	ABA24728
ORF7	386	Oxidoreductase	Putative aldo/keto reductase (<i>Tricodesmium erythraeum</i> IMS101)	70%/84% (383)	ZP00328942
ORF8	339	Sulfotransferase	Putative sulfotransferase (<i>Shewanella</i> sp. PV-4)	22%/39% (228)	EAP04083
ORF9	664	ABC-type transporter	NcpC (<i>Nostoc</i> sp. ATCC 53789)	57%/75% (663)	AAO23332
ORF10	257	Succinate dehydrogenase	Succinate dehydrogenase (<i>Sulfolobus tokodaii</i>)	33%/53% (238)	BAB40685

Finally, AerF and BacC, respectively, show close similarity to dehydrogenases and reductases. On the basis of the preceding analysis, we propose that AerC, AerD, AerE, and AerF provide the Choi moiety of aeruginoside. A biochemical model for the biosynthetic route to Choi is discussed below.

Downstream of the *aerD-F* region, the *aerG* gene encodes a NRPS module with the domain order C, A, PCP, C, and PCP. The analysis of the substrate binding pocket of the A domain revealed closest similarity to enzymes activating proline or methylproline, although part of the specificity conferring residues differ from typical proline-activating domains. This suggests that a proline-like amino acid rather than proline itself is activated (Table S1). It is interesting to note that spumigin, a peptide isolated from a *Nodularia* strain, is similar to aeruginosin but contains a methylproline moiety instead of Choi [35]. In *Planktothrix*, however, it is expected that AerG directly activates and incorporates the Choi moiety (see also feeding studies below). It is not clear to which extent the second C domain and, in particular, the second PCP domain of AerG are involved in aeruginoside biosynthesis. Amide bond forma-

tion principally functions with a free-standing C domain, such as for the VibH enzyme in vibrobactin biosynthesis [36]. The second C and PCP domains of AerG in *Planktothrix* may be remnants of a complete NRPS module. It is of note that aeruginosin-type metabolites are known from *Microcystis aeruginosa* in which the C-terminal carboxyl group of arginine is reduced to the aldehyde and/or alcohol oxidation state [4, 6]. Production of metabolites of the latter type requires a functional, complete module and a reductase in order to effect release of the peptide from the enzyme. The structure of AerG from the *Planktothrix aer* cluster suggests that in this organism the synthetase may be able to add the agmatine/Aeap moiety to the growing peptide chain with just a C domain. Since agmatine and Aeap lack the carboxyl that would otherwise tether the growing peptide to the synthetase, a thioesterase or domain for a reductive release is not required in the *Planktothrix* enzyme. We therefore postulate that the release of the peptide from the enzyme complex is triggered by amide bond formation between the PCP-bound carboxyl group of Choi and the primary amine of agmatine or of Aeap.




Figure 3. Biosynthetic Pathway of Aeruginosides 126A and 126B

(A) Schematic representation of the aeruginoside biosynthesis gene cluster of *Planktothrix agardhii* CYA126/8. Genes encoding NRPS or NRPS/PKS hybrid components are indicated in light and dark gray, respectively. Genes presumably involved in Choi biosynthesis are checked, and tailoring genes are highlighted with stripes. Genes that were not associated with aeruginoside biosynthesis are shown in white.

(B) Model for the formation of aeruginoside A and B and predicted domains of AerA-AerI. Each circle represents a NRPS domain or tailoring function. A, adenylation domain; KR, ketoreductase domain; ACP, acyl carrier protein; C, condensation domain; PCP, peptidyl carrier protein; E, epimerization domain.

The AerH protein shows similarity to oxygenases including isopenicillin N synthase and related enzymes from various bacteria. This enzyme may have a function in Aeap biosynthesis from arginine or agmatine. However, since the origin of carbon atoms in the Aeap residue is not presently known, this assignment is speculative at this time. The putative tailoring enzyme AerI shows similarity to glycosyltransferases. A glycosyltransferase is needed for the transfer of the xylose moiety to the hydroxyl group of Choi. Although the specificity of the glycosyltransferase cannot be deduced from the sequence, the close proximity of the gene to aeruginosin biosynthesis genes suggests a role in aeruginoside biosynthesis.

Taken together, the in silico analysis of the enzymes encoded by the *aer* gene cluster suggests roles for the enzymes in aeruginoside biosynthesis. The majority of enzymatic reactions proposed are in agreement with the synthesis of the tetrapeptide (Figure 3B). However, several of the enzymatic steps remain obscure at this point, particularly as far as Aeap biosynthesis is concerned. In order to obtain further insights into some of the reactions involved in Choi biosynthesis, we constructed additional mutants and initiated feeding studies.

Mutagenesis of Putative Choi Biosynthesis Genes and Feeding Studies

In order to confirm the proposed involvement of AerD, AerE, and AerF in the synthesis of the Choi moiety, two knockout constructs were designed. For both constructs,

a chloramphenicol resistance cassette was inserted into the *aerD*–*aerF* region. The first construct was bearing a deletion in the *aerE*–*aerF* region, whereas in the second experiment the antibiotic cassette was inserted into the *aerD* gene (Figure 4A). Both constructs were introduced into *Planktothrix* by electroporation. Mutants obtained after 3 months were analyzed by PCR and revealed the successful homologous recombination of the construct DNA (Figure 4B). Extracts of broths from both mutants, designated Δ *aerEF* and Δ *aerD*, were compared to the wild-type by LC-ESI-MS (Figures 4C, 4D, and 4F). While the wild-type spectrum revealed the characteristic aeruginoside peak at 10 min (m/z 714.80–715.80), no aeruginoside could be detected in the mutants (Figures 4C, 4D, and 4F). Following this observation, both mutants were fed with external L-Choi hydrochloride [37] and reanalyzed by LC-ESI-MS. Mass spectra of cell mass of the mutant after Choi feeding clearly revealed the production of aeruginoside (Figures 4E and 4G). These results confirmed the proposed involvement of the *aerD*–*aerF* genes in Choi biosynthesis. Moreover, the results indicate that Choi is directly activated at the AerG enzyme rather than being synthesized as an enzyme-bound intermediate. This conclusion is further supported by the fact that free Choi could be detected in the wild-type strain *Planktothrix* CYA126/8 and in the Δ *aerB* mutant strain (data not shown).

Although AerD–AerF look strikingly similar to the BacA–BacC enzymes of *Bacillus*, the Choi moiety and the




Figure 4. Analysis of Putative Choi Biosynthesis Genes

(A and B) Inactivation of putative Choi biosynthesis genes by homologous recombination. (A) Schematic representation of $\Delta aerEF$ and $\Delta aerD$ knockout plasmids for the insertional mutagenesis of *P. agardhii* CYA126/8. Chloramphenicol resistance gene cassettes (Cmr) are highlighted in black. (B) PCR amplification with the DNA from wild-type (lanes 1 and 4), $\Delta aerEF$ and $\Delta aerD$ mutants (lanes 2 and 5), and $\Delta aerEF$ and $\Delta aerD$ plasmid constructs (lanes 3 and 6) with primers amplifying the *aerD*–*aerF* (lanes 1–3) and the *aerD* (lanes 4–6) regions, respectively. (C–G) LC-ESI MS (positive SIM, range *m/z* 714.80–715.80) of extracts from (C) *P. agardhii* CYA126/8 WT, (D) *P. agardhii* CYA126/8 $\Delta AerEF$, (E) *P. agardhii* CYA126/8 $\Delta AerEF$ + Choi, (F) *P. agardhii* CYA126/8 $\Delta AerD$, (G) *P. agardhii* CYA126/8 $\Delta AerD$ + Choi. 1; aeruginoside 126A.

anticapsin moiety exhibit only limited structural similarity. An adaptation of the enzymatic reactions of the bacilysin pathway to Choi biosynthesis is not straightforward as the sequence of reactions in *Bacillus* is not well understood. Both pathways have in common that direct feeding with tyrosine itself did not result in any evidence for its incorporation into either Choi (this work and data not shown) or anticapsin [38]. This suggests that a tyrosine precursor is the obligatory intermediate in these two pathways. In the case of anticapsin biosynthesis it has been suggested that prephenate, an intermediate of phenylalanine biosynthesis, might fulfill this function [33]. This view is supported by the observation that there is strong sequence similarity

between BacA and AerD and prephenate decarboxylases. However, although Choi bears an amino group, an enzyme potentially transferring an amino group is not encoded within the aeruginoside gene cluster. Therefore, we propose that the amino group of Choi is provided by arogenate (pretyrosine), a known intermediate of tyrosine biosynthesis in cyanobacteria [39]. Arogenate could potentially also serve as precursor for anticapsin biosynthesis (Figure 5A). On the basis of these assumptions, the hypothetical Choi biosynthetic pathway is proposed as follows (Figure 5B): arogenate is oxidized by AerC to produce **3**, which is cyclized via Michael addition. This step must be enzyme catalyzed in order for the correct




Figure 5. Proposed Biosynthetic Pathways of the Anticapsin Moiety of Bacilysin and the Choi Moiety of Aeruginosin

(A) Biosynthetic model for the formation of anticapsin starting from arogenate. (B) Biosynthetic model for the formation of Choi starting from arogenate.

diastereomer of Choi to be produced: addition of the α -amino group to the *pro-R* carbon of the dienone results in the 2*S*, 3*aR*, 7*aR* diastereomer of **4**, whereas addition to the *pro-S* carbon results in the 2*S*, 3*aS*, 7*aS* diastereomer of **4**. The resulting vinylogous β -keto acid **4** is decarboxylated by AerD to yield the bicyclic amino acid **5**, which may be reduced by AerF (Figure 5B). This model is chemically reasonable but will require additional refinement through future enzymological studies after the complete heterologous expression of the AerD–AerF region or through further mutagenesis studies.

SIGNIFICANCE

Many cyanobacterial metabolites are infamous as health-threatening toxins that occur in bacterial blooms. However, various natural products from these organisms show important biological activities, such as the aeruginosins, which specifically inhibit serine proteases and are regarded as promising drug candidates. Aeruginosins are also interesting from a structural point of view, as they represent unusual cyanobacterial peptides that contain a unique 2-carboxy-6-hydroxyoctahydroindole (Choi) moiety. We have identified a putative aeruginoside biosynthesis genes in the genome of the freshwater cyanobacterium *Planktothrix agardhii* CYA126 and succeeded in unequivocally proving its identity by directed mutagenesis of a NRPS core component encoded by the cluster. Comparison of the metabolic profiles led to the discovery and structural elucidation of two previously unreported glycopeptides, which were named aeruginosides 126A and 126B. The aeruginoside biosynthesis pathway from *Planktothrix* CYA126/8 provides a novel example for a complex molecular assembly line composed of NRPS featuring an unusual loading module as well as an extender module for the unique Choi moiety and a putative glycosyltransferase. Heterologous expression of aeruginoside pathway genes for sustainable production or engineering of potential aeruginoside-derived drugs would be to be a promising alternative to chemical de novo synthesis in the future.

EXPERIMENTAL PROCEDURES

Bacterial Strains and Culturing

The axenic strain *P. agardhii* CYA126/8 was provided by Prof. K. Sivonen (University of Helsinki, Helsinki, Finland). Wild-type and mutant strains were cultured in Z8 medium as described previously [16]. Cultures were maintained under continuous white light ($30 \mu\text{E m}^{-2} \text{s}^{-1}$) on a shaker with 40 or 80 rpm at 23°C. DNA manipulations were performed in *Escherichia coli* XL1 Blue in LB medium. *E. coli* XL1 Blue MRF⁺ Kan strain was used for the excision of lambda phagemids, according to the suggestions of the manufacturer (Stratagene, La Jolla, CA).

Construction and Screening of the Phagemid Library

The isolation of chromosomal DNA from *P. agardhii* CYA126/8 and the construction of a lambda ZAP library were previously reported [16].

Screening of the library was performed by hybridization. Clone gaps were connected by PCR or inverse PCR, respectively, by using *Pfu* polymerase (Stratagene, La Jolla, CA).

Analysis of Aeruginosides

Cells of WT strain CYA126/8 and the mutant cultures grown in BG₁₁ medium²² at 20°C ($15 \mu\text{E m}^{-2} \text{s}^{-1}$) were harvested on preweighed glass fiber filters (GF/C). The cell mass was freeze dried, and the filters were reweighed for the determination of the dry weight (6 mg of DW each for WT and the mutant). Filters were extracted in aqueous MeOH (50/50, v/v) on a shaker for 60 min. The extract was centrifuged, and the supernatant cleaned up by solid phase extraction (SPE) with Sep-Pak tC18 cartridges (Waters, Vienna, Austria). The tC18 cartridges were conditioned by using MeOH followed by equilibration with 25% aq. MeOH (v/v). Samples were applied in 25% aq. MeOH, the cartridge was washed with five bed volumes of the same solvent, and aeruginosins were eluted with 80% MeOH (v/v). The cleaned extract was resuspended in 100 μl of 50% MeOH, and 100 μl were injected with HPLC-DAD (HP1100) and separated on a LiChrospher 100, ODS, 5 μm , LiChroCART 250-4 cartridge system (Merck, Darmstadt, Germany) by using a linear gradient of aq. acetonitrile (with 0.05% v/v trifluoroacetic acid, TFA) starting with 20% acetonitrile (v/v) and increasing up to 50% acetonitrile within 45 min with a flow of 1 ml min^{-1} . Peptides were detected at 210 nm.

Isolation of Aeruginosides

Forty-five liters of CYA126/8 were grown as described above, harvested by filtration through a sieve with a mesh size of 3 μm by using low pressure vacuum, and the cell mass was freeze dried. Cells were extracted with 50% MeOH (see above), and the extracts injected after SPE purification as described. HPLC purification was performed by using a linear gradient of aqueous acetonitrile (0.05% v/v TFA) starting with 20% acetonitrile (v/v) and increasing up to 30% acetonitrile within 15 min with a flow of 1 ml min^{-1} . The compounds were colorless oils and HPLC analysis of the purified compounds did not show UV absorbing contaminants.

Aeruginoside 126A (1)

Amorphous colorless solid; $[\alpha]_D^{25}$ 137 (c 0.0025 H₂O); ¹H NMR and ¹³C NMR, see Table 1; HRESITOFMS (positive ion mode) $[\text{M}+\text{H}]^+$ 715.4056 (715.4031 calculated).

Aeruginoside 126B (2)

Amorphous colorless solid; $[\alpha]_D^{25}$ 122 (c 0.0039, H₂O); ¹H NMR (500 MHz), DMSO-*d*₆ δ Plac: 7.25 (2H, t, *J* = 7.4 Hz, H-6/8), 7.21 (2H, d, *J* = 7.4 Hz, H-7), 7.17 (1H, m, H-5/9), 4.11 (1H, dd, *J* = 8.8, 3.7 Hz, H-2), 2.96 (1H, dd, *J* = 13.4, 3.7 Hz, H-3 α), 2.64 (1H, dd, *J* = 13.4, 8.8 Hz, H-3 β); Leu: 7.77 (1H, d, *J* = 7.8 Hz, NH), 4.53 (1H, m, H-2), 1.48 (1H, m, H-4), 1.46 (1H, m, H-3 α), 1.34 (1H, m, H-3 β), 0.87 (3H, d, *J* = 6.4 Hz, H-5), 0.84 (3H, d, *J* = 6.8 Hz, H-5'); Choi: 4.18 (1H, dd, *J* = 8.9, 8.5 Hz, H-2), 4.16 (1H, m, H-7a), 3.85 (1H, m, H-6), 2.30 (1H, m, H-3a), 2.29 (1H, m, H-7_{eq}), 2.09 (1H, m, H-4_{ax}), 2.03 (1H, m, H-3 α), 1.83 (1H, m, H-3 β), 1.61 (1H, m, H-7_{ax}), 1.57 (1H, brd, *J* = 12.7 Hz, H-5_{eq}), 1.50 (1H, m, H-5_{ax}), 1.48 (1H, m, H-4_{eq}); Agm: 7.74 (1H, t, *J* = 5.6 Hz, NH-1), 7.44 (1H, t, *J* = 5.6 Hz, NH-4), 3.08 (2H, m, H-4), 3.05 (2H, m, H-1), 1.44 (2H, m, H-2), 1.43 (2H, m, H-3); Xyl: 4.82 (1H, d, *J* = 3.6 Hz, H-1), 3.40 (1H, dd, *J* = 9.0, 9.0 Hz, H-3), 3.39 (1H, m, H-5_{eq}), 3.34 (1H, dd, *J* = 11.0, 11.0 Hz, H-5_{ax}), 3.28 (1H, m, H-4), 3.22 (1H, dd, *J* = 9.0, 3.6 Hz, H-2); ¹³C NMR (125 MHz, DMSO-*d*₆) δ Plac: 173.2 ((1) C-1), 138.5 ((1) C-4), 129.4 ((2) C-5/9), 128.0 ((2) C-6/8), 126.0 ((2) C-7), 72.06 ((2) C-2), 40.46 ((3) C-3); Leu: 169.8 ((1) C-1), 48.5 ((2) C-2), 41.3 ((3) C-3), 24.2 ((2) C-4), 23.4 ((4) C-5'), 21.9 ((4) C-5); Choi: 171.6 ((1) C-1), 70.12 ((2) C-6), 59.9 ((2) C-2), 54.1 ((2) C-7a), 36.1 ((2) C-3a), 30.5 ((3) C-3), 29.3 ((3) C-7), 24.5 ((3) C-5), 19.5 ((3) C-4); Agm: 156.7 ((1) C-5), 40.56 ((3) C-4), 37.9 ((3) C-1), 26.2 ((3) C-2), 25.8 ((3) C-3); Xyl: 96.6 ((2) C-1), 73.5 ((2) C-3), 72.0 ((2) C-2), 70.07 ((2) C-4), 62.2 ((3) C-5); HRESITOFMS (positive ion mode) $[\text{M}+\text{H}]^+$ 691.4051 (691.4031 calculated, Δ = 2.0 mmu).

Analysis of Stereochemistry of 1

Aeruginoside A (0.5 mg) was heated in 0.5 ml 6 N HCl at 110°C for 16 hr. After cooling to rt, the solution was extracted with EtOAc (3 × 0.5 ml), and the organic layer was removed under a stream of nitrogen. The residue was dissolved in MeOH, and a portion was injected onto a Chiralpak MA(+) column equilibrated in 2 mM aq. CuSO₄/CH₃CN (85:15) and eluted at 0.8 ml min⁻¹. D-Plac eluted at 41 min (retention time L-Plac: 59 min).

The aqueous layer from the hydrolysis was evaporated under a stream of nitrogen, and a portion of the residue was injected onto a Chirex D column equilibrated in 2 mM aq. CuSO₄/CH₃CN (95:5, v/v) and eluted at 0.8 ml min⁻¹. D-Leu eluted at 61.7 min (retention time L-Leu: 51.4 min).

Protease Inhibition Assays

Serine protease inhibitory activities were analyzed as described previously [40] by using porcine pancreas trypsin (Sigma; St. Louis, USA), bovine plasma thrombin (Sigma; St. Louis, USA), and human plasma plasmin (Sigma; St. Louis, USA). Aeruginosin 98A from *M. aeruginosa* NIES98 was used as a positive control.

Construction of the Gene Disruption Plasmids for aerB, aerD, and aerEF

The phagemid pBK-CMV-ΔaerB was digested with *Xcm*I (E1 site in *emp*erization domain). After blunting of the phagemide ends by Klenow treatment, the 1.4 kb *Bsa*AI fragment from pACYC184 containing the chloramphenicol resistance cassette was inserted. Positive clones containing the antibiotic resistance cassette were identified by restriction analysis. The phagemid pBK-CMV-ΔaerD was digested with *Eco*RI and *Xho*I to remove a *Hind*III-recognition site. Subsequently, sticky ends of the plasmid were blunted as described above. The resulting plasmid was purified (Qiaquick, QIAGEN, Hilden, Germany) and self-ligated. The derived plasmid was digested with *Hind*III and then blunted as described above. The 1.4 kb *Bsa*AI fragment from pACYC184 containing the chloramphenicol resistance cassette was cloned into the blunted plasmid. Positive clones and the orientation of the antibiotic cassette were identified by restriction analysis.

A 6115 bp gene fragment containing *aerEF* was amplified by PCR with the primers Pa36_61Arv3 (5'-AAG CAG GAA GCT TCA CTT GG-3') and Dhg1641rv5 (5'-CCT GCA ATC ACA AAC AAT TAC T-3') with *Pfu* polymerase (Stratagene, La Jolla, CA). The resulting PCR product was treated with *Taq* polymerase (72°C for 30 min) and purified with a gel purification kit (Qiaquick, QIAGEN, Hilden, Germany). The purified DNA was cloned into the pDrive vector (QIAGEN, Hilden, Germany). A 1,178 bp fragment covering the region between 5' end of *aerE* to the mid of *aerF* was deleted with *Pac*I and blunted by Klenow treatment. The 1.4 kb *Bsa*AI fragment from pACYC184 containing the chloramphenicol resistance cassette was cloned into the blunted plasmid generating the gene disruption plasmid pDrive-ΔaerEF. The antibiotic resistance clones were identified by restriction analysis.

Planktothrix Mutagenesis

The three constructs (15 to 20 μg of plasmid DNA) were linearized in the multicloning site of the vectors using *Bam*HI. The restricted DNA was column purified (Qiaquick, QIAGEN, Hilden, Germany), denaturated at 95°C for 10 min, and then immediately put onto an ice bath. Fifty milliliters of cultured algal cells were centrifuged at 6,000 rpm and washed three times with 1 mM HEPES buffer and centrifuged. Cells were resuspended in 1 mM HEPES and subjected to electroporation with 10 μg of the single-strand plasmid DNA (1.0 kV, 25 μF, 200 Ω).

Transformed cyanobacterial cells were inoculated in 100 ml of Z8 medium and cultivated as described above. After three days, 50 μg of chloramphenicol was added to culture medium. After 4 to 6 weeks, a fresh Z8 medium containing 0.5 μg ml⁻¹ chloramphenicol was added to the cultures. Transformants were purified by increasing stepwise the chloramphenicol concentration up to 5 μg ml⁻¹ (8 to 12 weeks).

DNA Sequencing and Sequence Analysis

Sequencing of phagemids was performed by primer walking with the Big Dye terminator cycle sequencing kit (ABI, Foster City, USA). Analyses of DNA sequences and deduced amino acid sequences were performed with the NCBI (National Institute for Biotechnology Information, Bethesda, MD) BLAST server (<http://www.ncbi.nlm.nih.gov/>) and the ExpASY proteomics server (<http://www.expasy.ch/>) of the Swiss Institute of Bioinformatics (Geneve, Switzerland).

HPLC-ESIMS Analysis of Aeruginosides and Feeding Experiments

One 200 ml aliquot of wild-type and two 200 ml aliquots of the ΔaerD and ΔaerEF mutant cells were cultivated as described above. One aliquot of the ΔaerD and ΔaerEF mutant, respectively, was fed with 2 mg L-Choi•HCl in the early logarithmic phase. All cultures were grown until they reached late logarithmic phase (OD750: 1.0), centrifuged, and freeze dried.

Freeze-dried algal cells were extracted twice with 80% MeOH (10 ml/100 ml cultured cells) and once with MeOH (10 ml/100 ml cultured cells). Combined 80% MeOH and MeOH extracts were concentrated to an aqueous suspension, which was then washed with diethylether. The aqueous layer was extracted with *n*-BuOH. The *n*-BuOH layer was evaporated under reduced pressure and dissolved in 50% MeOH (0.5 ml/100 ml cultured cells). Sample solutions were analyzed by LC-ESIMS (Finnigan TSQ Quantum ULTRA ThermoFinnigan Surveyor HPLC system, Thermo Electron Corporation, Waltham, MA, USA) by using reversed-phase HPLC (PRONTOSIL 120-5-C18 ace-EPS (4.0 × 250 mm, Bischoff Chromatography, Leonberg, Germany) with a linear gradient of 10% to 80% acetonitrile containing 0.1% AcOH in 35 min at a flow rate of 1.0 ml min⁻¹.

Supplemental Data

Supplemental Data include a comparison of the putative specificity-conferring codes of adenylation domains involved in aeruginoside biosynthesis, NMR spectra of aeruginosides, and 1D NOE and ROESY correlations for the Choi residue and are available at <http://www.chembiol.com/cgi/content/full/14/5/565/DC1/>.

ACKNOWLEDGMENTS

K.I. was financially supported by Japan Society for Promotion of Science Postdoctoral Fellowships for Research abroad. We thank Dr. O. Scheibner and Mrs. A. Perner for LC-ESI MS analyses and Dr. M. Ito-Ishida for assistance. The work was supported by a grant of the EU (PEPCY) to T.B. and E.D. and by the National Science Foundation (OCE04-32479) and the National Institute of Environmental Health Sciences (P50 ES012740) to T.H.

Received: December 13, 2006

Revised: April 2, 2007

Accepted: April 3, 2007

Published: May 29, 2007

REFERENCES

1. Welker, M., and von Döhren, H. (2006). Cyanobacterial peptides—nature's own combinatorial biosynthesis. *FEMS Microbiol. Rev.* 30, 530–563.
2. Murakami, M., Okita, Y., Matsuda, H., Okino, T., and Yamaguchi, K. (1994). Aeruginosin 298-A, a thrombin and trypsin-inhibitor from the blue-green-alga *Microcystis-aeruginosa* (NIES-298). *Tetraedron Lett.* 35, 3129–3132.
3. Kodani, S., Ishida, K., and Murakami, M. (1998). Aeruginosin 103-A, a thrombin inhibitor from the cyanobacterium *Microcystis viridis*. *J. Nat. Prod.* 61, 1046–1048.
4. Matsuda, H., Okino, T., Murakami, M., and Yamaguchi, K. (1996). Aeruginosins 102-A and B, new thrombin inhibitors from the

- cyanobacterium *Microcystis viridis* (NIES-102). *Tetrahedron* 52, 14501–14506.
- Murakami, M., Ishida, K., Okino, T., Okita, Y., Matsuda, H., and Yamaguchi, K. (1995). Aeruginosin-98-A and aeruginosin-98-B, trypsin-inhibitors from the blue-green-alga *Microcystis aeruginosa* (NIES-98). *Tetrahedron Lett.* 36, 2785–2788.
 - Ishida, K., Okita, Y., Matsuda, H., Okino, T., and Murakami, M. (1999). Aeruginosins, protease inhibitors from the cyanobacterium *Microcystis aeruginosa*. *Tetrahedron* 55, 10971–10988.
 - Shin, H.J., Matsuda, H., Murakami, M., and Yamaguchi, K. (1997). Aeruginosins 205A and -B, serine protease inhibitory glycopeptides from the cyanobacterium *Oscillatoria agardhii* (NIES-205). *J. Org. Chem.* 62, 1810–1813.
 - Banker, R., and Carmeli, S. (1999). Inhibitors of serine proteases from a waterbloom of the cyanobacterium *Microcystis* sp. *Tetrahedron* 55, 10835–10844.
 - Ploutno, A., Shoshan, M., and Carmeli, S. (2002). Three novel protease inhibitors from a natural bloom of the cyanobacterium *Microcystis aeruginosa*. *J. Nat. Prod.* 65, 973–978.
 - Carroll, A.R., Buchanan, M.S., Edser, A., Hyde, E., Simpson, M., and Quinn, R.J. (2004). Dysinosins B-D, inhibitors of factor VIIa and thrombin from the Australian sponge *Lamellodysidea chlorea*. *J. Nat. Prod.* 67, 1291–1294.
 - Carroll, A.R., Pierens, G.K., Fechner, G., de Almeida Leone, P., Ngo, A., Simpson, M., Hyde, E., Hooper, J.N.A., Bostrom, S.L., Musil, D., and Quinn, R.J. (2002). Dysinosin A: a novel inhibitor of factor VIIa and thrombin from a new genus and species of Australian sponge of the family dysideidae. *J. Am. Chem. Soc.* 124, 13340–13341.
 - Sandler, B., Murakami, M., and Clardy, J. (1998). Atomic structure of the trypsin-aeruginosin 98-B complex. *J. Am. Chem. Soc.* 120, 595–596.
 - Steiner, J.L.R., Murakami, M., and Tulinsky, A. (1998). Structure of thrombin inhibited by aeruginosin 298-A from a blue-green alga. *J. Am. Chem. Soc.* 120, 597–598.
 - Dittmann, E., Neilan, B.A., and Börner, T. (2001). Molecular biology of peptide and polyketide biosynthesis in cyanobacteria. *Appl. Microbiol. Biotechnol.* 57, 467–473.
 - Schwarzer, D., Finking, R., and Marahiel, M.A. (2003). Nonribosomal peptides: from genes to products. *Nat. Prod. Rep.* 20, 275–287.
 - Christiansen, G., Fastner, J., Erhard, M., Börner, T., and Dittmann, E. (2003). Microcystin biosynthesis in *Planktothrix*: genes, evolution, and manipulation. *J. Bacteriol.* 185, 564–572.
 - Nishizawa, T., Ueda, A., Asayama, M., Fujii, K., Harada, K., Ochi, K., and Shirai, M. (2000). Polyketide synthase gene coupled to the peptide synthetase module involved in the biosynthesis of the cyclic heptapeptide microcystin. *J. Biochem. (Tokyo)* 127, 779–789.
 - Rouhiainen, L., Vakkilainen, T., Siemer, B.L., Buikema, W., Haselkorn, R., and Sivonen, K. (2004). Genes coding for hepatotoxic heptapeptides (microcystins) in the cyanobacterium *Anabaena* strain 90. *Appl. Environ. Microbiol.* 70, 686–692.
 - Tillett, D., Dittmann, E., Erhard, M., von Döhren, H., Börner, T., and Neilan, B.A. (2000). Structural organization of microcystin biosynthesis in *Microcystis aeruginosa* PCC7806: an integrated peptide-polyketide synthetase system. *Chem. Biol.* 7, 753–764.
 - Chang, Z.X., Flatt, P., Gerwick, W.H., Nguyen, V.A., Willis, C.L., and Sherman, D.H. (2002). The barbamide biosynthetic gene cluster: a novel marine cyanobacterial system of mixed polyketide synthase (PKS)-non-ribosomal peptide synthetase (NRPS) origin involving an unusual trichloroleucyl starter unit. *Gene* 296, 235–247.
 - Chang, Z.X., Sitachitta, N., Rossi, J.V., Roberts, M.A., Flatt, P.M., Jia, J.Y., Sherman, D.H., and Gerwick, W.H. (2004). Biosynthetic pathway and gene cluster analysis of curacin A, an antitubulin natural product from the tropical marine cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* 67, 1356–1367.
 - Edwards, D.J., Marquez, B.L., Nogle, L.M., McPhail, K., Goeger, D.E., Roberts, M.A., and Gerwick, W.H. (2004). Structure and biosynthesis of the jamaicamides, new mixed polyketide-peptide neurotoxins from the marine cyanobacterium *Lyngbya majuscula*. *Chem. Biol.* 11, 817–833.
 - Rouhiainen, L., Paulin, L., Suomalainen, S., Hyytiäinen, H., Buikema, W., Haselkorn, R., and Sivonen, K. (2000). Genes encoding synthetases of cyclic depsipeptides, anabaenopeptilides, in *Anabaena* strain 90. *Mol. Microbiol.* 37, 156–167.
 - Edwards, D.J., and Gerwick, W.H. (2004). Lyngbyatoxin biosynthesis: sequence of biosynthetic gene cluster and identification of a novel aromatic prenyltransferase. *J. Am. Chem. Soc.* 126, 11432–11433.
 - Stachelhaus, T., Mootz, H.D., and Marahiel, M.A. (1999). The specificity-conferring code of adenylation domains in nonribosomal peptide synthetases. *Chem. Biol.* 6, 493–505.
 - Fujii, K., Sivonen, K., Adachi, K., Noguchi, K., Shimizu, Y., Sano, H., Hirayama, M., Suzuki, M., and Harada, K. (1997). Comparative study of toxic and non-toxic cyanobacterial products: a novel glycoside, suomilide, from non-toxic *Nodularia spumigena* HKVV. *Tetrahedron Lett.* 38, 5529–5532.
 - Ploutno, A., and Carmeli, S. (2005). Banyasin and banyasides A and B, three novel modified peptides from a water bloom of the cyanobacterium *Nostoc* sp. *Tetrahedron* 61, 575–583.
 - Valls, N., Vallribera, M., Font-Bardia, M., Solans, X., and Bonjoch, J. (2003). Synthesis of the proposed core of aeruginosins 205: the new alpha -amino acid (2S,3aS,6R,7aS)-2-carboxy-6-chlorooctahydroindole. *Tetrahedron Asymmetry* 14, 1241–1244.
 - Hicks, L.M., Moffitt, M.C., Beer, L.L., Moore, B.S., and Kelleher, N.L. (2006). Structural characterization of in vitro and in vivo intermediates on the loading module of microcystin synthetase. *ACS Chem. Biol.* 1, 93–102.
 - Moore, B.S., and Hertweck, C. (2002). Biosynthesis and attachment of novel bacterial polyketide synthase starter units. *Nat. Prod. Rep.* 19, 70–99.
 - Steinborn, G., Hajirezaei, M.R., and Hofemeister, J. (2005). bac genes for recombinant bacilysin and anticapsin production in *Bacillus* host strains. *Arch. Microbiol.* 183, 71–79.
 - Kunst, F., Ogasawara, N., Moszer, I., Albertini, A.M., Alloni, G., Azevedo, V., Bertero, M.G., Bessieres, P., Bolotin, A., Borchert, S., et al. (1997). The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* 390, 249–256.
 - Hilton, M.D., Alaeddinoglu, N.G., and Demain, A.L. (1988). Synthesis of bacilysin by *Bacillus subtilis* branches from prephenate of the aromatic amino acid pathway. *J. Bacteriol.* 170, 482–484.
 - McLuskey, K., Cameron, S., Hammerschmidt, F., and Hunter, W.N. (2005). Structure and reactivity of hydroxypropylphosphonic acid epoxidase in fosfomicin biosynthesis by a cation- and flavin-dependent mechanism. *Proc. Natl. Acad. Sci. USA* 102, 14221–14226.
 - Fujii, K., Sivonen, K., Adachi, K., Noguchi, K., Sano, H., Hirayama, K., Suzuki, M., and Harada, K. (1997). Comparative study of toxic and non-toxic cyanobacterial products: novel peptides from toxic *Nodularia spumigena* AV1. *Tetrahedron Lett.* 38, 5525–5528.
 - Keating, T.A., Marshall, C.G., Walsh, C.T., and Keating, A.E. (2002). The structure of VibH represents nonribosomal peptide synthetase condensation, cyclization and epimerization domains. *Nat. Struct. Biol.* 9, 522–526.

37. Valls, N., Vallibera, M., Lopez-Canet, M., and Bonjoch, J. (2002). Synthesis of microcin SF608. *J. Org. Chem.* *67*, 4945–4950.
38. Roscoe, J., and Abraham, E.P. (1966). Experiments relating to biosynthesis of bacilysin. *Biochem. J.* *99*, 793–800.
39. Hall, G.C., Flick, M.B., Gherna, R.L., and Jensen, R.A. (1982). Biochemical diversity for biosynthesis of aromatic-amino-acids among the cyanobacteria. *J. Bacteriol.* *149*, 65–78.
40. Shin, H.J., Murakami, M., Matsuda, H., and Yamaguchi, K. (1996). Microviridins D-F, serine protease inhibitors from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *Tetrahedron* *52*, 8159–8168.

Accession Numbers

The sequence of the *aer* gene cluster and flanking regions has been deposited in the EMBL database under the accession number AM071396.