

## Refereed journal papers and book chapters (May 2016)

- 91 **Weisse, T.** & Sonntag, B. (2016). Ciliates in planktonic food webs: communication and adaptive response. In: Witzany, G. & Nowacki, M. (eds), *Biocommunication of ciliates*, Spinger, Berlin (in press)
- 90 **Weisse, T.**, Gröschl, B. & Kremser, V. (2016). Phytoplankton response to short-term temperature and nutrient changes. *Limnologica* (accepted 02/05/16)
- 89 Cieplinski, A., **Weisse, T.** & Obertegger, U. (2016). High diversity in *Keratella cochlearis* (Rotifera, Monogononta) - morphological and genetic evidence. *Hydrobiologia* (accepted 21/03/2016/)
- 88 Weckström, K., Weckström, J., Huber, K., Kamenik, C., Schmidt, R., Salvenmoser, W., Rieradevall, M., **Weisse, T.**, Psenner, R & Kurmayer, R. (2016). Impacts of climate warming on Alpine lake biota over the last decade. *Arctic, Antarctic, and Alpine Research* 48, 361-376.
- 87 **Weisse, T.**, Anderson, R., Arndt, H., Calbet, A., Hansen, P.J. & Montagnes, D.J.S. (2016). Functional Ecology of aquatic phagotrophic protists – concepts, limitations, and perspectives. *Eur. J. Protistol.*, <http://dx.doi.org/10.1016/j.ejop.2016.03.003>.
- 86 Dunthorn, M., Lipps, J.H., Dolan, J.R., Abboud-Abi Saab, M., Aescht, E., Bachy, C., Barría de Cao, M.S., Berger, H., Bourland, W.A., Choi, J.K., Clamp, J., Doherty, M., Gao, F., Gentekaki, E., Gong, J., Hu, X., Huang, J., Kamiyama, T., Johnson, M.D., Kammerlander, B., Kim, S.Y., Kim, Y.-O., la Terza, A., Laval-Peuto, M., Lipscomb, D., Lobban, C.S. Long, H., Luporini, P., Lynn, D.H., Macek, M., Mansergh, R.I., Martín-Cereceda, M., McManus, G.G., Montagnes, D.J.S., Ong'ondo, G.O., Patterson, D.J., Pérez-Uz, B., Quintela-Alonso, P., Safi, L.S.L., Santoferrara, L.F. Sonntag, B., Song, W., Stoeck, T., Stoecker, D.K., Strüder-Kypke, M.C., Trautmann, I., Utz, L.R.P., Vallesi, A., Vd'ačný, P., Warren, A., **Weisse, T.**, Wickham, S.A., Yi, Z., Zhang, W. Zhan, Z., Zufall, R. & Agatha, S. (2015). Ciliates – protists with complex morphologies and ambiguous early fossil record. *Mar. Micropaleontol.* 119, 1–6.
- 85 Nevalainen, L., Luoto, T.P., Manca, M. & **Weisse, T.** (2015). A paleolimnological perspective on aquatic biodiversity in Austrian mountain lakes. *Aquat. Sci.* 77:59–69 [DOI 10.1007/s00027-014-0363-6].
- 84 **Weisse, T.** (2014). Ciliates and the Rare Biosphere – Community Ecology and Population Dynamics. *J. Eukaryot. Microbiol.* 61, 419–433.
- 83 **Weisse, T.**, Scheffel, U., Stadler, P. & Foissner, W. (2013d). Functional ecology of the ciliate *Glaucomides bromelicola*, and comparison with the sympatric species *Bromeliothrix metopoides*. *J. Eukaryot. Microbiol.* 60, 578-587.
- 82 **Weisse, T.**, Scheffel, U., Stadler, P. & Foissner, W. (2013c). *Bromeliothrix metopoides*, a boom and bust ciliate (Ciliophora, Colpodea) from tank bromeliads. *Eur. J. Protistol.* 49, 406-419 [doi: 10.1016/j.ejop.2013.02.001].
- 81 **Weisse, T.**, Laufenstein, N. & Weithoff, G. (2013b). Multiple environmental stressors confine the ecological niche of the rotifer *Cephalodella acidophila*. *Freshwater Biol.* 58: 1008-1015 [doi:10.1111/fwb.12104].
- 80 **Weisse, T.**, Moser, M., Scheffel, U., Stadler, P., Berendonk, T., Weithoff, G. & Berger, H. (2013a). Systematics and species-specific response to pH of *Oxytricha acidotolerans* sp. nov. and *Urosomoida* sp. (Ciliophora, Hypotricha) from acid mining lakes. *Eur. J. Protistol.* 49, 255-271, [doi: 10.1016/j.ejop.2012.08.001].
- 79 **Weisse, T.**, Berendonk, T., Kamjunke, N., Moser, M., Scheffel, U., Stadler, P. & Weithoff, G. (2011). Significant habitat effects influence protist fitness: evidence for local adaptation from acidic mining lakes. *Ecosphere* 2: art134 [doi:http://dx.doi.org/10.1890/ES11-00157.1].
- 78 Jersabek, C., Weithoff, G. & **Weisse, T.** (2011). *Cephalodella acidophila* n. sp. (Monogononta: Notommatidae), a new rotifer species from highly acidic mining lakes. *Zootaxa* 2939: 50-58.
- 77 Moser, M. & **Weisse, T.** (2011c). The most acidified Austrian lake in comparison to a neutralized mining lake. *Limnologica* 41: 303-315.
- 76 Moser, M. & **Weisse, T.** (2011b). Combined stress effect of pH and temperature narrows the niche width of flagellates in acid mining lakes. *J. Plankton Res.* 33: 1023-1032.
- 75 Moser, M. & **Weisse, T.** (2011a). The outcome of competition between the two

- chrysoomonads *Ochromonas* sp. and *Poterioochromonas malhamensis* depends on pH. *Eur. J. Protistol.* 47: 79-85.
- 74 Weithoff, G., Moser, M.; Kamjunke, N., Gaedke, U. & **Weisse, T.** (2010): Lake morphometry strongly shapes the plankton community structure in acidic mining lakes. *Limnologia* 40: 161-166.
- 73 Moser, M., Callieri, C. & **Weisse, T.** (2009). Photosynthetic and growth response of freshwater picocyanobacteria are strain-specific and sensitive to photoacclimation. *J. Plankton Res.* 31: 349-357.
- 72 **Weisse, T.**, Boenigk, J. & Müller, H. (2009). Ecological methods for the study of heterotrophic nano- and microplankton of fresh and marine waters. In: A course in Protozoology, Röttger, R., Knight, R. & Foissner, W. (eds), *Protozool. Monographs*: 4: 232-242.
- 71 Montagnes, D.J.S., Morgan, G., Bissinger, J.E., Atkinson, D. & **Weisse, T.** (2008). Short-term temperature change may impact freshwater carbon flux: a microbial perspective. *Global Change Biology* 14: 2810-2822.
- 70 **Weisse, T.**, Strüder-Kypke, M., Berger, H. & Foissner, W. (2008). Genetic, morphological, and ecological diversity of spatially separated clones of *Meseres corlissi* Petz and Foissner, 1992 (Ciliophora, Spirotrichea). *J. Eukaryot. Microbiol.* 55: 257-270.
- 69 Gächter, E. & **Weisse, T.** (2008). Long-term acclimation of growth rates in the oligotrich freshwater ciliate *Meseres corlissi*. *Verh. Internat. Verein. Limnol.* 30: 218-222.
- 68 **Weisse, T.** (2008) Distribution and diversity of aquatic protists: an evolutionary and ecological perspective. *Biodivers. Conserv.* 17:243-259.
- 67 **Weisse, T.**, Scheffel, U., Stadler, P. & Foissner, W. (2007). Local adaptation among geographically distant clones of the cosmopolitan freshwater ciliate *Meseres corlissi*. II. Response to pH. *Aquat. Microb. Ecol.* 47: 289-297.
- 66 **Weisse, T.** (2006). Biodiversity of freshwater microorganisms - achievements, problems, and perspectives. In: GLIWICZ, Z. M., MAZURKIEWICZ-BORON, G. & ROUEN, K. (eds.): *Advances in European Freshwater Sciences, 2005. Pol. J. Ecol.* 54: 633-652.
- 65 **Weisse, T.** (2006) Biodiversity of aquatic protists – what can we learn from bacteria and *Daphnia*? *Endocytobiosis Cell Res.* 17: 154-163.
- 64 Gächter, E. & **Weisse, T.** (2006). Local adaptation among geographically distant clones of the cosmopolitan freshwater ciliate *Meseres corlissi*. I. Temperature response. *Aquat. Microb. Ecol.* 45: 291-300.
- 63 Foissner, W., Pichler, M., Al-Rahsheid, K. & **Weisse, T.** (2006) The unusual, Lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichea): encystment and genesis and release of the lepidosomes. *Acta Protozool.* 45: 323-338.
- 62 **Weisse, T.** (2006). Freshwater ciliates as ecophysiological model organisms - lessons from *Daphnia*, major achievements, and future perspectives. *Arch. Hydrobiol.* 167:371-402.
- 61 **Weisse, T.** & Stadler, P. (2006) Effect of pH on growth, cell volume, and production of freshwater ciliates, and implications for their distribution. *Limnol. Oceanogr.* 51: 1708-1715.
- 60 Müller, H., Foissner, W. & **Weisse, T.** (2006) The role of soil in the life cycle of *Meseres corlissi* (Ciliophora: Oligotrichea): experiments with two clonal strains from the type locality, an astatic meadow pond. *Aquat. Microb. Ecol.* 42: 199-208.
- 59 **Weisse, T.** & Rammer, S. (2006). Pronounced ecophysiological clonal differences of two common freshwater ciliates, *Coleps spetai* (Prostomatida) and *Rimostrombidium lacustris* (Oligotrichida), challenge the morphospecies concept. *J. Plankton Res.* 28: 55-63.
- 58 Foissner, W., Müller, H. & **Weisse, T.** (2005). The unusual, Lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichea): Light and scanning electron microscopy, cytochemistry. *Acta Protozool.* 44: 201-215.
- 57 Callieri, C., Moro, S., Crosbie, N.D. & **Weisse, T.** (2005) Strain-specific photosynthetic response of freshwater picocyanobacteria. *Verh. int. Ver. Limnol.* 29: 777-782.
- 56 **Weisse, T.** (2004). *Meseres corlissi*: a rare oligotrich ciliate adapted to warm water and temporary habitats. *Aquat. Microb. Ecol.* 37: 75-83.
- 55 **Weisse, T.** (2003). Pelagic Microbes - Protozoa and the Microbial Food Web. In: O'Sullivan, P. & Reynolds, C.S. (eds.), *The Lakes Handbook, Vol. I*, Blackwell Scientific Publ., Oxford, pp. 417-460.
- 54 Crosbie, N.D., Pöckl, M. & **Weisse, T.** (2003). Rapid establishment of clonal isolates of freshwater autotrophic picoplankton by single-cell and single-colony sorting. *J. Microbiol. Meth.* 55: 361-370.
- 53 Crosbie, N.D., Pöckl, M. & **Weisse, T.** (2003). Dispersal and phylogenetic diversity of non-marine picocyanobacteria, inferred from

- 16SrRNA gene and *cpcBA*-intergenic spacer sequence analyses. *Appl. Environ. Microbiol.* 69: 5716-5721.
- 52 Crosbie, N.D., Teubner, K. & **Weisse, T.** (2003). Flow-cytometric mapping provides novel insights into the seasonal and vertical distribution of freshwater autotrophic picoplankton. *Aquat. Microb. Ecol.* 33: 53-66.
- 51 Lindström, E.S., Stadler, P. & **Weisse, T.** (2003). Live sorting and survival of unstained and DAPI-stained ciliates by flow cytometry. *Arch. Hydrobiol.* 157: 173-184.
- 50 **Weisse, T.** & Mindl, B. (2002). Picocyanobacteria - sensitive bioindicators of contaminant stress in an alpine lake (Lake Traunsee, Austria). *Water, Air, and Soil Pollution: Focus* 2: 191-210
- 49 Müller, H., Stadler, P. & **Weisse, T.** (2002). Seasonal dynamics of cyst formation of strombidid ciliates in alpine Lake Mondsee, Austria. *Aquat. Microb. Ecol.* 29: 181-188.
- 48 **Weisse, T.** (2002) The significance of inter- and intraspecific variation in bacterivorous and herbivorous protists. *Antonie van Leeuwenhoek* 81: 327-341.
- 47 **Weisse, T.**, Stadler, P., Lindström, E.S., Kimmanse, S.A. & Montagnes, D.J.S. (2002). Interactive effect of temperature and food concentration on growth rate: A test case using the small freshwater ciliate *Urotricha farcta*. *Limnol. Oceanogr.* 47: 1447-1455.
- 46 **Weisse, T.**, Gomoiu, M.-T., Scheffel, U. & Brodrecht, F. (2002). Biomass and size composition of the combjelly *Mnemiopsis* sp. in the north-western Black Sea during spring and summer. *Estuar. Cst. Shelf Sci.* 54: 423-437.
- 45 Lindström, E.S., **Weisse, T.** & Stadler, P. (2002). Enumeration of small ciliates in culture by flow cytometry and nucleic acid staining. *J. Microb. Methods* 49: 173-182.
- 44 **Weisse, T.** & Frahm, A. (2002). Direct and indirect impact of two common rotifer species (*Keratella* spp.) on two abundant ciliate species (*Urotricha furcata*, *Balanion planctonicum*). *Freshwater Biology* 47: 53-64.
- 43 **Weisse, T.**, Karstens, N., Meyer, V.C.M., Janke, L. Lettner, S. & Teichgräber, K (2001). Niche separation in common prostome freshwater ciliates: the effect of food and temperature. *Aquat. Microb. Ecol.* 26: 167-179.
- 42 **Weisse, T.** & Frahm, A. (2001). Species-specific interactions between small planktonic ciliates (*Urotricha* spp.) and rotifers (*Keratella* spp.). *J. Plankton Res.* 23: 1329-1338.
- 41 Clevén, E.-J. & **Weisse, T.** (2001). Seasonal succession and taxon-specific bacterial grazing rates of heterotrophic nanoflagellates in Lake Constance. *Aquat. Microb. Ecol.* 23: 147-161.
- 40 Munawar, M., Munawar, I.F., Dermott, R., Munawar, S.F., Norwood, W., Lynn, D., **Weisse, T.**, Johannssen, O.E., Wenghofer, C., Niblock, H., Carou, S. & Gasenbeck, K. (2001). Aquatic Ecosystem Health of Fathom Five National Marine Park: A structural and functional assessment. In: Parker, S. & Munawar, M. (eds.), Ecology, Culture, and Conservation of a protected Area: Fathom Five National Marine Park, pp. 99-149. Backhuys Publ., Leiden, The Netherlands.
- 39 Montagnes, D.J.S. & **Weisse, T.** (2000): Fluctuating temperatures affect growth and production rates of planktonic ciliates. *Aquat. Microb. Ecol.* 21: 97-102.
- 38 **Weisse, T.** & Gomoiu, M.-T. (2000). Biomass and size structure of the moon jellyfish *Aurelia aurita* during spring and summer. *J. Plankton Res.* 22: 223-239.
- 37 **Weisse, T.** & McIsaac, E.A. (2000). Significance and fate of bacterial production in oligotrophic lakes in British Columbia. *Can. J. Fish. Aquat. Sci.* 57: 96-105.
- 36 **Weisse, T.** (1999). Bacterivory in the northwestern Indian Ocean during the intermonsoon - NE monsoon period. *Deep-Sea Res.* II, 46: 795-814.
- 35 Munawar, M., Munawar, I.F., **Weisse, T.**, van Stam, H., Fitzpatrick, M., Lorimer, J., Wenghofer, C. & Carou, S. (1999). Probing microbial food web structure in Lake Erie. In: Munawar, M., Edsall, T. & Munawar, I.F. (eds.), *State of Lake Erie (SOLE): Past, Present and Future*, pp. 417-439. Ecovision World Monograph Series, Backhuys Publ., Leiden, The Netherlands.
- 34 **Weisse, T.** & Müller, H. (1998). Planktonic protozoa and the microbial food web. *Arch. Hydrobiol. Spec. Issues Advanc. Limnol.* 53: 223-254.
- 33 Gaedke, U. & **Weisse, T.** (1998). Seasonal and interannual variability of picocyanobacteria in Lake Constance (1987-1997). *Arch. Hydrobiol. Spec. Issues Advanc. Limnol.* 53: 143-158.
- 32 Sterner, R.W., Clasen, J., Lampert, W. & **Weisse, T.** (1998). Carbon: phosphorus stoichiometry and food chain production. *Ecology Letters* 1: 146-151.
- 31 **Weisse, T.** & Montagnes, D.J.S. (1998). The effect of temperature on inter- and intraspecific isolates of *Urotricha* (Prostomatida, Ciliophora). *Aquat. Microb. Ecol.* 15: 285-291.
- 30 **Weisse, T.** (1997) Growth and production of heterotrophic nanoflagellates in a mesoeutrophic lake. *J. Plankton Res.* 19: 703-722.

- 29 **Weisse, T.** & Kirchoff, B. (1997). Feeding of the heterotrophic dinoflagellate *Peridiniopsis berlinense* on cryptophytes: analysis by flow cytometry and electronic particle counting *Aquat. Microb. Ecol.* 12: 153-164.
- 28 Zimmermann, U., Müller, H. & **Weisse, T.** (1996). Seasonal and spatial variability of planktonic heliozoa in Lake Constance. *Aquat. Microb. Ecol.* 11: 21-29.
- 27 Munawar, M., Munawar, I.F. & **Weisse, T.** (1995b). Utility and applicability of autotrophic picoplankton as bioindicators of ecosystem health. In: Munawar, M., Hanninen, O., Roy, N., Munawar, I.F., Karenlampi, L. & Brown, L. (eds.), *Bioindicators of Environmental Health*, pp. 13-27. SPB Academic Publishing, Amsterdam, The Netherlands.
- 26 Müller, H. & **Weisse, T.** (1994). Laboratory and field observations on the scuticociliate *Histiobalantium* from the pelagic zone of Lake Constance, FRG. *J. Plankton Res.* 16: 391-401.
- 25 **Weisse, T.**, Tande, K., Verity, P., Hansen, F. & Gieskes, W. (1994). The trophic significance of *Phaeocystis* blooms. In Lancelot, C. & Wassmann, P. (eds.), *Ecology of Phaeocystis dominated ecosystems*. *J. Mar. Syst.* 5: 67-79.
- 24 **Weisse, T.** (1994). Structure of microbial food webs in relation to the trophic status of lakes and fish grazing pressure: a key role of cyanobacteria? In: Pinto-Coelho, R.M., Giani, A., and Sperling, E.v. (eds), *Ecology and Human Impact on Lakes and Reservoirs in Minas Gerais*, pp. 55-70, SEGRAC, Brazil.
- 23 Munawar, M., Munawar, I.F., **Weisse, T.**, Lepard, G.G. & Legner, M. (1994). The significance and future potential of using microbes for assessing ecosystem health: The Great Lakes example. *J. Ecosyst. Health* 3: 295-30.
- 22 **Weisse, T.** & Stockner, J.G. (1993). Eutrophication: The role of microbial food webs. *Mem. Ist. ital. Idrobiol.* 52: 133-150.
- 21 **Weisse, T.** (1993). Dynamics of autotrophic picoplankton in freshwater and marine ecosystems. In: Jones, J.G. (ed.), *Adv. Microb. Ecol.* 13: 327-370.
- 20 Gradinger, R., **Weisse, T.** & Pillen, T. (1992). Significance of picocyanobacteria in the Red Sea and the Gulf of Aden. *Bot. Mar.* 35: 245-250.
- 19 **Weisse, T.** (1991b). The annual cycle of heterotrophic freshwater nanoflagellates: role of bottom-up versus top-down control. *J. Plankton Res.* 13: 167-185.
- 18 **Weisse, T.** (1991a). The microbial loop and its sensitivity to eutrophication and contaminant enrichment: a cross-system overview. *Int. Revue ges. Hydrobiol.* 76: 327-337.
- 17 **Weisse, T.** & Scheffel-Möser, U. (1991). Uncoupling the microbial loop: growth and grazing loss rates of bacteria and heterotrophic nanoflagellates in the North Atlantic. *Mar. Ecol. Prog. Ser.* 71: 195-205.
- 16 Müller, H., Schöne, A., Pinto-Coelho, R.M., Schweizer, A. & **Weisse, T.** (1991). Seasonal succession of ciliates in Lake Constance. *Microb. Ecol.* 21: 119-138.
- 15 **Weisse, T.** & Kenter, U. (1991). Ecological characteristics of autotrophic picoplankton in a prealpine lake. *Int. Revue ges. Hydrobiol.* 76: 493-504.
- 14 **Weisse, T.**, Müller, H., Pinto-Coelho, R.M., Schweizer, A., Springmann, D. & Baldringer, G. (1990). Response of the microbial loop to the phytoplankton spring bloom in a large prealpine lake. *Limnol. Oceanogr.* 35: 781-794.
- 13 **Weisse, T.** & Scheffel-Möser, U. (1990b). Growth and grazing loss rates in single-celled *Phaeocystis* cf. *pouchetii* (Prymnesiophyceae). *Mar. Biol.* 106: 153-158.
- 12 **Weisse, T.** & Scheffel-Möser, U. (1990a). Morphometric characteristics and carbon content of *Phaeocystis* cf. *pouchetii*. *Bot. Mar.* 33: 197-203.
- 11 **Weisse, T.** & Müller, H. (1990). Significance of heterotrophic nanoflagellates and ciliates in large lakes: Evidence from Lake Constance. In Tilzer, M.M. & Serruya, C. (eds.), *Large lakes - ecological structure and function*, pp. 540-555. Springer/Science Tech. Publ., Madison.
- 10 **Weisse, T.** (1990). Trophic interactions among heterotrophic microplankton, nano-plankton, and bacteria in Lake Constance. *Hydrobiologia* 191: 111-122.
- 9 **Weisse, T.** (1989). The microbial loop in the Red Sea: dynamics of pelagic bacteria and heterotrophic nanoflagellates. *Mar. Ecol. Prog. Ser.* 55: 241-250.
- 8 **Weisse, T.** & Rudstam, L. (1989). Excretion and respiration rates of *Neomysis integer* (Mysidaceae): effects of temperature, sex and starvation. *Hydrobiologia* 178: 253-258.
- 7 Munawar, M. & **Weisse, T.** (1989). Is the 'microbial loop' an early warning indicator of anthropogenic stress? *Hydrobiologia* 188/189: 163-174.
- 6 **Weisse, T.** (1988). Dynamics of autotrophic picoplankton in Lake Constance. *J. Plankton Res.* 10: 1179-1188.

- 5 Lancelot, C., Billen, G., Sournia, A., **Weisse, T.**, Colijn, F., Veldhuis, M.J.W., Davis, A. & Wassmann, P. (1987). *Phaeocystis* blooms and nutrient enrichment in the continental coastal zones of the North Sea. *Ambio* 16: 38-46.
- 4 **Weisse, T.**, Grimm, N., Hickel, W. & Martens, P. (1986). Dynamics of *Phaeocystis pouchetii* blooms in the Wadden Sea of Sylt (German Bight, North Sea). *Estuar. Cstl. Shelf Sci.* 23: 171-182.
- 3 Grimm, N. & **Weisse, T.** (1985). Die Temperaturabhängigkeit des Wachstums von *Phaeocystis pouchetii* (Haptophyceae) in Batchkulturen. *Helgoländer Meeresunters.* 39: 201-211 (in German with English abstract).
- 2 Schneider, G. & **Weisse, T.** (1985). Metabolism measurements of *Aurelia aurita* planulae larvae, and calculation of maximal survival period of the free swimming stage. *Helgoländer Meeresunters.* 39: 43-47.
- 1 **Weisse, T.** (1983). Feeding of calanoid copepods in relation to *Phaeocystis pouchetii* blooms in the German Wadden Sea area off Sylt. *Mar. Biol.* 74: 87-94.
- Weisse, T.** (2008b). Wilhelm FOISSNER and the German Society for Protozoology. *Denisia* 23: 9.
- Weisse, T.** (2008a). Limnoecology - The Ecology of Lakes and Streams (Book review). *J. Plankton Res.* 30: 489-490.
- Weisse, T.** (2007b) Limnology in Austria and at Mondsee - Going Against the Trend? *SILnews* 51: 7-8.
- Weisse, T.** (2007a). Der Artbegriff bei Bakterien, Ein- und Vielzellern. *Denisia* 20: 149-154.
- Weisse, T.** (2003). YURI SOROKIN: The Black Sea-Ecology and Oceanography. *Int. Rev. ges. Hydrobiol.* 88: 427-428.
- Weisse, T.** & Lettner, S. (2002). The ecological significance of intraspecific variation among freshwater ciliates. *Verh. int. Ver. Limnol.* 28: 1880-1884.
- Weisse, T.**, Springmann, D.B., Müller, H. (1995). Ökologie des heterotrophen Nanoplanktons im Süßwasser und Meer. In: Röttger, R. (ed.), *Praktikum der Protozoologie*, pp. 40-47. G. Fischer Vlg., Stuttgart.
- Geller, W., Berberovic, R., Gaedke, U., Müller, H., Pauli, H.-R., Tilzer, M.M. & **Weisse, T.** (1991). Relations among the components of autotrophic and heterotrophic plankton during the seasonal cycle 1987 in Lake Constance. *Verh. int. Ver. Limnol.* 24: 831-836.
- Weisse, T.** & Schweizer, A. (1991). Seasonal and interannual variation of autotrophic picoplankton in a large prealpine lake (Lake Constance). *Verh. int. Ver. Limnol.* 24: 821-825.

### Conference Proceedings, non-refereed papers, and book reviews

- Weisse, T.** (2014) Acidic Pit Lakes – The Legacy of Coal and Metal Surface Mines; W. Geller, M. Schultze, R. Kleinmann, C. Wolkersdorfer (Eds.). *Limnologica* 48: 80-81.
- Weisse, T.** (2008c) The meaning of protist diversity: ecology meets taxonomy. *Denisia* 23: 297-306.

### Contributions as editor

- Weisse, T.** (2013). Editorial. *Eur. J. Protistol.* 49 (3): 327.
- Weisse, T.** (2012). Editorial. *Eur. J. Protistol.* 48 (2): 95.
- Weisse, T.** (2010). Editorial. *Eur. J. Protistol.* 46 (4): 253.
- Weisse, T.** (2009). The European Journal of Protistology – Changes, chances and challenges (Editorial). *Eur. J. Protistol.* 45: 163–164.
- Simon, M., Güde, H. & **Weisse, T.** (eds.) (1996). Aquatic Microbial Ecology, Proceedings of the Sixth International Workshop on the Measurement of Microbial Activities in the Cycling of Matter in Aquatic Environments, Konstanz, Germany, 17-22 April 1995. *Arch. Hydrobiol. Spec. Issues Advanc. Limnol.* 48.