

Yegor Dikarev

Dear ISEM Team! Here are my solutions for the 1st lecture.

ex. 1. Prove that $\sin(nx), n \in \mathbb{N}$, form a complete orthogonal system in $L^2(0, \pi)$, compute the L^2 norms.

Proof. Let $f_n(x) = \sin(nx)$, then consider scalar product in $L^2(0, \pi)$ as

$$\begin{aligned} (f_n, f_m) &= \int_0^\pi f_n(x) \overline{f_m(x)} dx = \int_0^\pi \sin(nx) \sin(mx) dx = \frac{1}{2} \int_0^\pi (\cos(n-m)x - \cos(n+m)x) dx = \\ &= \frac{1}{2} \left(\int_0^\pi \cos(n-m)x dx - \int_0^\pi \cos(n+m)x dx \right) = \frac{1}{2} \left(\frac{\sin(n-m)x}{n-m} \Big|_0^\pi - \frac{\sin(n+m)x}{n+m} \Big|_0^\pi \right) = 0. \end{aligned}$$

Thus, $f_n(x) = \sin(nx)$ is orthogonal system, and it is complete because system $f_n(x) = \sqrt{\frac{2}{\pi}} \sin(nx)$ is dense in $L^2(0, \pi)$ (see 1.1 in lection 1) Computing the L^2 norms.

$$\begin{aligned} \|f_n\|_2 &= \left(\int_0^\pi \sin^2(nx) dx \right)^{\frac{1}{2}} = \left(\int_0^\pi \frac{1 - \cos(2nx)}{2} dx \right)^{\frac{1}{2}} = \\ &= \left(\frac{x}{2} \Big|_0^\pi - \frac{1}{2} \int_0^\pi \cos(2nx) dx \right)^{\frac{1}{2}} = \left(\frac{\pi}{2} - \frac{1}{4} \sin(2nx) \Big|_0^\pi \right)^{\frac{1}{2}} = \sqrt{\frac{\pi}{2}} \end{aligned}$$

ex. 3. Let X be a Banach space and $A_1 : X \rightarrow X$ and $A_2 : X \rightarrow X$ linear maps such that
1. $D(A_1) \subset D(A_2)$ and A_1 is a restriction of A_2 . 2. A_1 is surjective and A_2 is injective. Show that $A_1 = A_2$.

Proof. To prove that the operators are equivalent, we have to show that their domains are equal. According to Property 1, we have to show that $D(A_2) \subset D(A_1)$. We have $D(A_1) \subset D(A_2)$, consequently, because of the linearity of operators $Im A_1 \subset Im A_2$. A_1 is surjective, therefore $Im A_1 = X$ and $Im A_2 = X$. Let $x \in D(A_2) \setminus D(A_1)$, then $A_2 x = A_1 x_0, x_0 \in D(A_1)$ (A_1 is surjective), but $A_1 x_0 = A_2 x_0$, then $A_2 x = A_2 x_0$ and $A_2(x - x_0) = 0$, hence $x = x_0 \in D(A_2) \setminus D(A_1)$. Thus, $D(A_2) \subset D(A_1)$ and $D(A_2) = D(A_1)$.

ex. 5. Let $p \in [1, \infty)$ and consider the Banach space $L^p(\mathbb{R})$. Prove that the formula

$$(S(t)f)(x) = f(t+x) \text{ for } f \in L^p, x \in \mathbb{R}, t \geq 0$$

defines a strongly continuous semigroup on L^p . What happens for $p = \infty$?

Proof. To prove that $S(t)$ defines a strongly continuous semigroup, we have to check the properties from the semigroup definition.

1. $(S(t+s)f)(x) = f(x+t+s) = f((x+s)+t) = (S(t)f)(x+s) = (S(s)(S(t)f))(x) = (S(t)S(s)f)(x)$.
2. $(S(0)f)(x) = f(x+0) = f(x)$.
3. $S(f)$ is continuous, because $f \in BUC(\mathbb{R})$ (see 1.2 in Lecture 1).