

On the structure of universal differentiability sets.

Michael Dymond

Universität Innsbruck

Relations between Banach Space Theory and
Geometric Measure Theory,
University of Warwick,
Wednesday 10th June 2015.

Sets containing a point of differentiability of every Lipschitz function.

A set $S \subseteq \mathbb{R}^d$ is called a *universal differentiability set* if S contains a point of differentiability of every Lipschitz function $f : \mathbb{R}^d \rightarrow \mathbb{R}$.

A set $E \subseteq \mathbb{R}^d$ is called a *non-universal differentiability set* if there exists a Lipschitz function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ such that f is nowhere differentiable in E .

Examples.

- By Rademacher's theorem, any subset of \mathbb{R}^d of positive Lebesgue measure is a universal differentiability set.
- Universal differentiability sets in \mathbb{R} are precisely the sets of positive Lebesgue measure.

Existence of exceptional universal differentiability sets.

- (Preiss 1990) For $d \geq 2$, there exist UDSs in \mathbb{R}^d with Lebesgue measure zero.
- (Doré, Maleva 2011) For $d \geq 1$, \mathbb{R}^d contains compact UDSs with Hausdorff dimension one.
- (D., Maleva 2014) For $d \geq 1$, \mathbb{R}^d contains compact UDSs with Minkowski dimension one.
- (Preiss, Speight 2014) (Alberti, Csörnyei, Preiss 2010) (Csörnyei, Jones) \mathbb{R}^d contains Lebesgue null UDSs for Lipschitz mappings $f : \mathbb{R}^d \rightarrow \mathbb{R}^l$ if and only if $l < d$.

Connection to porosity.

Definition

A subset P of \mathbb{R}^d is called porous if there exists $c \in (0, 1)$ such that for every $x \in P$ and every $\epsilon > 0$ there exists $h \in \mathbb{R}^d$ with $\|h - x\| \leq \epsilon$ and $B(h, c \|h - x\|) \cap P = \emptyset$. P is called σ -porous if P can be expressed as a countable union of porous sets.

- Every porous subset of \mathbb{R}^d is a *non-universal differentiability set*.
- (Kirchheim, Preiss, Zajíček, 2001) Every σ -porous subset of \mathbb{R}^d is a *non-universal differentiability set*.

Can a universal differentiability set be decomposed?

Given a universal differentiability set $S \subseteq \mathbb{R}^d$, is it possible to write $S = A \cup B$ where A and B are non-universal differentiability sets?

Equivalently: Given a universal differentiability set $S \subseteq \mathbb{R}^d$ is it possible to find a pair of Lipschitz functions $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$ such that f and g have no common points of differentiability in S .

Simultaneous differentiability of Lipschitz functions.

- 1 (Lindenstrauss, Tišer, Preiss, 2012) Every pair (f, g) of Lipschitz functions on a Hilbert space have a common point of differentiability.
- 2 **Open Question:** Does every triple (f, g, h) of Lipschitz functions on a Hilbert space have a common point of differentiability?

Can a universal differentiability set be decomposed?

Does there exist a universal differentiability set $S \subseteq \mathbb{R}^d$ such that $S = A \cup B$ where A and B are non-universal differentiability sets?

Equivalently: Does there exist a universal differentiability set $S \subseteq \mathbb{R}^d$ for which it is possible to find a pair of Lipschitz functions $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$ such that f and g have no common points of differentiability in S .

The answer is yes.

- 1 (Csörnyei, Preiss, Tišer, 2004), (Alberti, Csörnyei, Preiss, 2010) Case $d = 2$.

Structural results for universal differentiability sets.

Theorem (D., 2014)

- 1 Let $S = A \cup B \subseteq \mathbb{R}^d$ be a universal differentiability set where A is a closed subset of S . Then either A or B is a universal differentiability set.
 - 2 Let $S = \bigcup_{i=1}^{\infty} A_i \subseteq \mathbb{R}^d$ be a universal differentiability set, where each A_i is a closed subset of S . Then at least one A_i is a universal differentiability set.
-
- Recall: (Csörnyei, Preiss, Tišer, 2004) There exists a universal differentiability set $S = A \cup B \subseteq \mathbb{R}^d$ such that both A and B are non-universal differentiability sets.

Structural results for universal differentiability sets.

Theorem (D., 2014)

- 1 Let $S = A \cup B \subseteq \mathbb{R}^d$ be a universal differentiability set where A is a closed subset of S . Then either A or B is a universal differentiability set.
 - 2 Let $S = \bigcup_{i=1}^{\infty} A_i \subseteq \mathbb{R}^d$ be a universal differentiability set, where each A_i is a closed subset of S . Then at least one A_i is a universal differentiability set.
-
- (Csörnyei, Preiss, Tišer, 2004) There exists a universal differentiability set $S = A \cup B \subseteq \mathbb{R}^d$ such that A is a G_δ set and both A and B are non-universal differentiability sets.

The kernel of a universal differentiability set.

Theorem (D., 2014)

Let S be a universal differentiability set and define

$$\ker(S) = S \setminus \{x \in S : \exists r > 0 \text{ s.t. } B(x, r) \cap S \text{ is a non-UDS}\}.$$

Then,

- 1 $\ker(S)$ is a universal differentiability set.
- 2 $\ker(\ker(S)) = \ker(S)$.

Open questions.

- 1 Does every universal differentiability set contain a closed universal differentiability set?
- 2 Does every subset of \mathbb{R}^d with positive Lebesgue measure contain a universal differentiability set of Lebesgue measure zero?

Thank you for listening.