

# Systems of coordinates in WLD Banach spaces

(Joint work with P. Hájek and T. Kania)

Tommaso Russo

tommaso.russo@unimi.it

*Università degli studi di Milano*

*Department of Mathematics 'Federigo Enriques'*

Joint meeting of  
UMI, SIMAI, PTM

Wrocław

September 17-20, 2018

## Systems of coordinates

- ▶ A collection  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma} \subseteq X \times X^*$  is a *biorthogonal system* if

$$\langle x_\alpha^*, x_\beta \rangle = \delta_{\alpha, \beta} \quad (\alpha, \beta \in \Gamma).$$

- ▶ If  $\|x_\gamma\| = \|x_\gamma^*\| = 1$ , the system is said to be an *Auerbach system*.
- ▶ An *M-basis* is a biorthogonal system such that:

$$\overline{\text{span}}\{x_\gamma\}_{\gamma \in \Gamma} = X \quad \& \quad \overline{\text{span}}^{w^*}\{x_\gamma^*\}_{\gamma \in \Gamma} = X^*.$$

- ▶ An *Auerbach basis* is an M-basis with  $\|x_\gamma\| = \|x_\gamma^*\| = 1$ .

**Auerbach (1934).** Every finite-dimensional normed space contains an Auerbach basis.

**Day (1962).** Every infinite-dimensional Banach space contains an infinite-dimensional subspace with an Auerbach basis, in particular it contains an infinite Auerbach system.

## And spaces with few of them

- ▶ **Kunen (1975).** (CH) There exists a non-separable Banach space that contains no uncountable biorthogonal system.
  - ▶ Other examples: (♣) Ostaszewski (1975), (◇) Shelah (1985).
  - ▶ **Todorčević (2006).** (MM) Every non-separable Banach space contains uncountable biorthogonal systems.
- ▶ **Johnson (1970).**  $\ell_\infty$  has no M-basis.
- ▶ **Plichko (1986).**  $c_0[0, 1] + C[0, 1]$  has no Auerbach basis.
- ▶ **Godun–Lin–Troyanski (1993).** Every non-separable Banach space  $X$  with  $B_{X^*}$   $w^*$ -separable, admits an equivalent norm with no Auerbach basis.
  - ▶ **Godun (1990).** The particular case  $X = \ell_1(c)$ .

**Problem (1):** Guirao, Montesinos, and Zizler (2016), *Open problems...*

Does there exist a non-separable Banach space  $X$  with unconditional basis such that no non-separable subspace of  $X$  has an Auerbach basis?

## 'Large' Banach spaces

### Theorem (Hájek, Kania, and R.)

*Let  $\kappa \geq \mathfrak{c}$  be a cardinal number and let  $X$  be a Banach space with  $w^*\text{-dens } X^* > \exp_2 \kappa$ . Then  $X$  admits a subspace  $Y$  with Auerbach basis and such that  $\text{dens } Y = \kappa^+$ .*

*Proof.* Set  $\lambda := w^*\text{-dens } X^*$ . By transfinite induction, find unit vectors  $(e_\alpha)_{\alpha < \lambda}$  and unit functionals  $(\varphi_{\alpha, \beta})_{\alpha < \beta < \lambda}$  such that:

- (i)  $\varphi_{\alpha, \beta}$  is a norming functional for  $e_\alpha - e_\beta \neq 0$  ( $\alpha < \beta < \lambda$ );
- (ii)  $e_\gamma \in \ker \varphi_{\alpha, \beta}$  ( $\alpha < \beta < \gamma < \lambda$ ).

We invoke the Erdős–Rado theorem, for the colouring

$$\{\alpha, \beta, \gamma\} \mapsto \langle \varphi_{\beta, \gamma}, e_\alpha \rangle \quad (\alpha < \beta < \gamma < \lambda).$$

We can build the desired Auerbach system out of a monochromatic set of cardinality  $\kappa^+$ .

## WLD spaces

A Banach space  $X$  is *weakly Lindelöf determined* (hereinafter, *WLD*) if the dual ball  $B_{X^*}$  is a Corson compact in the relative  $w^*$ -topology. For our purposes,  $X$  is WLD if it admits an M-basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  that *countably supports*  $X^*$ , i.e.,

$$\text{supp } x^* := \{\gamma \in \Gamma : \langle x^*, x_\gamma \rangle \neq 0\}$$

is a countable subset of  $\Gamma$ , for every  $x^* \in X^*$ .

### Theorem (Hájek, Kania, and R.)

*Every WLD Banach space  $X$  with  $\text{dens } X > \omega_1$  contains a subspace  $Y$  with Auerbach basis and such that  $\text{dens } Y = \text{dens } X$ .*

**Problem (2):** What happens if  $\text{dens } X = \omega_1$ ?

## The main result

### Theorem (Hájek, Kania, and R.)

(CH) *There exists a renorming  $\|\cdot\|$  of the space  $c_0(\omega_1)$  such that the space  $(c_0(\omega_1), \|\cdot\|)$  contains no uncountable Auerbach systems.*

Consequently, assuming the Continuum Hypothesis:

- ▶ There exists a non-separable Banach space with unconditional basis whose no non-separable subspace admits an Auerbach basis.
- ▶ There exists a WLD Banach space  $X$  with  $\text{dens } X = \omega_1$  every whose non-separable subspace fails to have an Auerbach basis.

Therefore, under CH, we can answer both Problems (1) and (2).

- ▶ For  $\alpha < \omega_1$ , let  $\sigma_\alpha$  be an enumeration of the set  $[0, \alpha)$ .
- ▶ Select  $\lambda \in (0, 1/6)$  and define  $\varphi_\alpha \in \ell_1(\omega_1)$  by

$$\varphi_\alpha(\eta) = \begin{cases} 1 & \text{if } \eta = \alpha \\ 0 & \text{if } \eta > \alpha \\ \lambda^k & \text{if } \eta < \alpha, \eta = \sigma_\alpha(k). \end{cases}$$

- ▶ Define a new (equivalent) norm on  $c_0(\omega_1)$  to be

$$\|x\| := \sup_{\alpha < \omega_1} |\langle \varphi_\alpha, x \rangle| \quad (x \in c_0(\omega_1)).$$

### The main property.

If  $u \in c_0(\omega_1) \setminus \{0\}$  and  $\text{supp } u < \alpha$ , then

$$\langle \varphi_\alpha, u \rangle := \sum_{\beta < \alpha} u(\beta) \langle \varphi_\alpha, e_\beta \rangle = \sum_{k=1}^{\infty} u(\sigma_\alpha(k)) \lambda^k$$

is a (non-trivial) real-analytic function of  $\lambda$  and therefore it has countably many zeros.

Consequently, for every countable subset  $(u_n)_{n=1}^{\infty}$  of  $c_0(\omega_1) \setminus \{0\}$ , we may select  $\lambda$  such that  $\langle \varphi_\alpha, u_n \rangle \neq 0$  for every  $n \in \mathbb{N}$ .

## Uncountable $(1+)$ -separated sets

A subset  $A$  of a normed space  $(X, \|\cdot\|)$  is  $(1+)$ -separated if  $\|x - y\| > 1$  for distinct  $x, y \in A$ .

### Theorem (T. Kania and T. Kochanek, 2016)

- ▶ Let  $X$  be a non-separable, reflexive Banach space. Then  $S_X$  contains an uncountable  $(1+)$ -separated subset;
- ▶ If  $K$  is a non-metrizable compact, the unit sphere of  $C(K)$  contains an uncountable  $(1+)$ -separated set.

Other relevant results are in the papers: Mercourakis–Vassiliadis 2015, Koszmider 2018, Cúth–Kurka–Vejnar 20--.



## Auerbach systems and (1+)-separation

### Theorem (Hájek, Kania, and R.)

*If  $X$  contains an Auerbach system with cardinality  $\mathfrak{c}^+$ , then the unit sphere of  $X$  contains an uncountable (1+)-separated subset.*

*Consequences:*

- ▶ Assume that  $w^*\text{-dens } X^* > \exp_2 \mathfrak{c}$ . Then both  $S_X$  and  $S_{X^*}$  contain an uncountable (1+)-separated subset.
- ▶ Let  $X$  be a WLD space with  $\text{dens } X > \mathfrak{c}$ . Then  $S_X$  and  $S_{X^*}$  contain uncountable (1+)-separated subsets.

**Note:** This is somewhat sharp, the conclusion can not be improved.

### Theorem (Hájek, Kania, and R.)

*Let  $\mathcal{F} \subseteq S_{c_0(\kappa)}$  be (1+)-separated. Then  $|\mathcal{F}| \leq \omega_1$ .*



**Thank you for your attention!**