



Czech Technical University in Prague

---

# Asplund Banach spaces and norming M-bases

Tommaso Russo

(Joint with P. Hájek, J. Somaglia, and S. Todorčević)

Institute of Mathematics  
Czech Academy of Sciences

June 23, 2020



- 1 Introduction and main results
- 2  $\varrho$ -functions and Theorem B
- 3 Proof of Theorem A
- 4 Semi-Eberlein compacta (*very* briefly)



- ① Introduction and main results
- ②  $\varrho$ -functions and Theorem B
- ③ Proof of Theorem A
- ④ Semi-Eberlein compacta (*very* briefly)



A system  $\{u_\alpha; \varphi_\alpha\}_{\alpha \in \Gamma} \subseteq \mathcal{X} \times \mathcal{X}^*$  is a **Markušević basis** (**M-basis**, for short) for  $\mathcal{X}$  if

- $\langle \varphi_\beta, u_\alpha \rangle = \delta_{\alpha, \beta}$ ,
- $\text{span}\{u_\alpha\}_{\alpha \in \Gamma}$  is dense in  $\mathcal{X}$ ,
- $\text{span}\{\varphi_\alpha\}_{\alpha \in \Gamma}$  is  $w^*$ -dense in  $\mathcal{X}^*$ .

$\{\langle \varphi_\alpha, x \rangle : \alpha \in \Gamma\}$  are the **coordinates** of  $x \in \mathcal{X}$

$\{\langle \psi, x_\alpha \rangle : \alpha \in \Gamma\}$  are the **coordinates** of  $\psi \in \mathcal{X}^*$ .

It is tempting to ask if  $\text{span}\{\varphi_\alpha\}_{\alpha \in \Gamma}$  exhausts  $\mathcal{X}^*$  in a stronger sense.

## Definition

An M-basis  $\{u_\alpha; \varphi_\alpha\}_{\alpha \in \Gamma}$  is  **$\lambda$ -norming** ( $0 < \lambda \leq 1$ ) if

$$\lambda \|x\| \leq \sup\{\langle \varphi, x \rangle : \varphi \in \text{span}\{\varphi_\alpha\}_{\alpha \in \Gamma}, \|\varphi\| \leq 1\}.$$

$\{u_\alpha; \varphi_\alpha\}_{\alpha \in \Gamma}$  is **shrinking** if  $\text{span}\{\varphi_\alpha\}_{\alpha \in \Gamma}$  is dense in  $\mathcal{X}^*$ .



- **Markušević, 1943.** Every separable Banach space has a 1-norming M-basis.
- **Amir–Lindenstrauss, 1968.** Every WCG Banach space has an M-basis;  
 $\implies$  Every reflexive space has a shrinking M-basis;  
*def* :  $\mathcal{X}$  is **WCG** if it contains a linearly dense weakly compact subset.
- Properties of M-bases can characterise classes of Banach spaces: weakly compact, shrinking, countably norming, ...
- What about norming M-bases?
- **John–Zizler, 1974.** Do WCG spaces have a norming M-basis?

## Theorem (Hájek, 2019)

There exists a WCG  $C(\mathcal{K})$  space with no norming M-basis.  
Actually,  $\mathcal{K}$  is uniform Eberlein, so  $C(\mathcal{K})$  is also Hilbert-generated.



# Towards the main result

*def* :  $\mathcal{X}$  is **Asplund** if every its separable subspace has separable dual.

- $C(\mathcal{K})$  is Asplund iff  $\mathcal{K}$  is scattered.
- **Troyanski, John–Zizler, Orihuela–Valdivia, Fabian, ...** TFAE:
  - (i)  $\mathcal{X}$  has a shrinking M-basis;
  - (ii)  $\mathcal{X}$  is WCG and Asplund;
  - (iii)  $\mathcal{X}$  is WLD and Asplund;
  - (iv)  $\mathcal{X}$  is WLD and  $\mathcal{X}^*$  admits a dual LUR norm.
- $\ell_1(\Gamma)$  admits a 1-norming M-basis.
  - Norming M-basis  $\not\Rightarrow$  WCG, or Asplund ( $\ell_1(\omega_1)$ );
  - Norming M-basis and WCG  $\not\Rightarrow$  Asplund ( $\ell_1$ ).
- **Godefroy ~1990.** Let  $\mathcal{X}$  be an Asplund space with a norming M-basis. Is  $\mathcal{X}$  WCG?

## Theorem A

There exists an Asplund space  $\mathcal{X}$  with a 1-norming M-basis such that  $\mathcal{X}$  is not WLD.



# A peculiar compact space

- $\mathcal{P}(\Gamma) \equiv \{0, 1\}^\Gamma$  by  $A \leftrightarrow 1_A$ ;
- This gives a compact ‘product’ topology on  $\mathcal{P}(\Gamma)$ ;
- If  $A \in \mathcal{P}(\Gamma)$ ,  $a_0, \dots, a_n \in A$ ,  $b_0, \dots, b_n \notin A$

$$\mathcal{U} := \{B \in \mathcal{P}(\Gamma) : a_0, \dots, a_n \in B, b_0, \dots, b_n \notin B\}$$

is a nghd of  $A$  in  $\mathcal{P}(\Gamma)$ .

## Theorem B

There exists a family  $\mathcal{F}_\omega \subseteq [\omega_1]^{<\omega}$  of finite subsets of  $\omega_1$  such that  $\mathcal{K}_\omega := \overline{\mathcal{F}_\omega}$  has the following properties:

- (i)  $\{\alpha\} \in \mathcal{K}_\omega$  for every  $\alpha < \omega_1$ ,
- (ii)  $[0, \alpha) \in \mathcal{K}_\omega$  for every  $\alpha \leq \omega_1$ ,
- (iii) if  $A \in \mathcal{K}_\omega$  is an infinite set, then  $A = [0, \alpha)$  for some  $\alpha \leq \omega_1$ ,
- (iv)  $\mathcal{K}_\omega$  is scattered.



- ① Introduction and main results
- ②  $\varrho$ -functions and Theorem B**
- ③ Proof of Theorem A
- ④ Semi-Eberlein compacta (*very* briefly)





**S. Todorčević, Partitioning pairs of countable ordinals, *Acta Math.* 159 (1987), 261–294.**



**S. Todorčević, *Walks on ordinals and their characteristics*. Birkhäuser Verlag, Basel, 2007.**

- We consider functions  $\varrho: [\omega_1]^2 \rightarrow \omega$ .
- We identify  $[\omega_1]^2 = \{(\alpha, \beta) \in \omega_1^2 : \alpha < \beta\}$ .
  - Thus, we write  $\varrho(\alpha, \beta)$ , with  $\alpha < \beta$ , for  $\varrho(\{\alpha, \beta\})$ .
- We also add the ‘boundary condition’  $\varrho(\alpha, \alpha) = 0$ .

## Definition (Todorčević)

A  **$\varrho$ -function** on  $\omega_1$  is a function  $\varrho: [\omega_1]^2 \rightarrow \omega$  such that:

( $\varrho 1$ )  $\{\xi \leq \alpha : \varrho(\xi, \alpha) \leq n\}$  is finite, for every  $\alpha < \omega_1$  and  $n < \omega$ ,

( $\varrho 2$ )  $\varrho(\alpha, \gamma) \leq \max\{\varrho(\alpha, \beta), \varrho(\beta, \gamma)\}$  for  $\alpha < \beta < \gamma < \omega_1$ ,

( $\varrho 3$ )  $\varrho(\alpha, \beta) \leq \max\{\varrho(\alpha, \gamma), \varrho(\beta, \gamma)\}$  for  $\alpha < \beta < \gamma < \omega_1$ .



## Proposition (Todorčević)

There exists a function  $\varrho: [\omega_1]^2 \rightarrow \omega$  such that  $(\alpha < \beta < \gamma < \omega_1)$ :

- $\varrho(\alpha, \beta) > 0$ ;
- $\varrho(\alpha, \gamma) \leq \max\{\varrho(\alpha, \beta), \varrho(\beta, \gamma)\}$ ;
- $\varrho(\alpha, \gamma) \neq \varrho(\beta, \gamma)$ .

$$F_n(\alpha) := \{\xi \leq \alpha : \varrho(\xi, \alpha) \leq n\}$$

$$\mathcal{F}_\varrho := \{F_n(\alpha) : n < \omega, \alpha < \omega_1\} \quad \text{and} \quad \mathcal{K}_\varrho := \overline{\mathcal{F}_\varrho}.$$

## Fact

- $F_0(\alpha) = \{\alpha\}$ ;
- $|F_n(\alpha)| \leq n + 1$ ;
- $(F_n(\alpha))_{n < \omega}$  converges to  $[0, \alpha]$ .

## Th B again, aka $\mathcal{K}_Q$ verifies Th B

The compact space  $\mathcal{K}_Q$  has the following properties:

- (i)  $\{\alpha\} \in \mathcal{K}_Q$  for every  $\alpha < \omega_1$ ,
- (ii)  $[0, \alpha) \in \mathcal{K}_Q$  for every  $\alpha \leq \omega_1$ ,
- (iii) if  $A \in \mathcal{K}_Q$  is an infinite set, then  $A = [0, \alpha)$  for some  $\alpha \leq \omega_1$ ,
- (iv)  $\mathcal{K}_Q$  is scattered.

**Proof.** (i)  $\{\alpha\} = F_0(\alpha) \in \mathcal{K}_Q$ . ✓

(ii) If  $\alpha = \alpha' + 1$ ,  $[0, \alpha) = [0, \alpha'] = \lim_{n < \omega} F_n(\alpha') \in \mathcal{K}_Q$ .

If  $\alpha$  is limit,  $([0, \beta + 1))_{\beta < \alpha}$  converges to  $[0, \alpha)$ . ✓

(iii)  $\implies$  (iv)

- Let  $\mathcal{D} \subseteq \mathcal{K}_Q$  be closed. Pick  $D_0 \in \mathcal{D}$  s.t.  $\exists \alpha < \beta$ ,  $\alpha \notin D_0$ ,  $\beta \in D_0$ .
- Pick a maximal element  $M \in \mathcal{D}$  with  $\alpha \notin M$ ,  $D_0 \subseteq M$ .
- $M$  is a finite set, by (iii).
- $\mathcal{U} := \{D \in \mathcal{D} : \alpha \notin D, M \subseteq D\} = \{M\}$ , so  $M$  is isolated in  $\mathcal{D}$ . ✓



(iii) if  $A \in \mathcal{K}_\varrho$  is infinite, then  $A = [0, \alpha)$  for some  $\alpha \leq \omega_1$ ;

def  $F_n(\alpha) := \{\xi \leq \alpha : \varrho(\xi, \alpha) \leq n\}$ .

Assume first  $|A| = \omega$ .

- Pick  $\alpha \in A$  and  $\tilde{\alpha} < \alpha$ ; we need  $\tilde{\alpha} \in A$ .
- Pick a sequence  $(F_{n_k}(\alpha_k))_{k < \omega} \rightarrow A$  (Fréchet–Urysohn property).
- If  $(n_k)_{k < \omega}$  is bounded, then  $|F_{n_k}(\alpha_k)| \leq M$ ; so,  $A$  is finite.
- WLOG,  $\varrho(\tilde{\alpha}, \alpha) \leq n_k$ .
- Also, assume  $\alpha \in F_{n_k}(\alpha_k)$ , namely  $\alpha \leq \alpha_k$  and  $\varrho(\alpha, \alpha_k) \leq n_k$ .
- By triangle inequality,

$$\varrho(\tilde{\alpha}, \alpha_k) \leq \max\{\varrho(\tilde{\alpha}, \alpha), \varrho(\alpha, \alpha_k)\} \leq n_k,$$

so  $\tilde{\alpha} \in F_{n_k}(\alpha_k)$ . Passing to the limit,  $\tilde{\alpha} \in A$ .  $\checkmark$



$\Sigma(\Gamma) := \{x \in [0, 1]^\Gamma : \text{supp}(x) \text{ is countable}\}.$

## Theorem (Dewille–Godefroy, Kalenda)

Let  $\mathcal{K} \subseteq [0, 1]^{\omega_1}$  be a compact set such that  $\mathcal{K} \cap \Sigma(\omega_1)$  is dense in  $\mathcal{K}$ . Let  $x \in \mathcal{K} \setminus \Sigma(\omega_1)$ . Then there exists an embedding  $\varphi: [0, \omega_1] \rightarrow \mathcal{K}$ :

- (i)  $\varphi(\alpha) \in \mathcal{K} \cap \Sigma(\omega_1)$ , for  $\alpha < \omega_1$ ,
- (ii)  $\text{supp}(\varphi(\alpha)) \subseteq \text{supp}(\varphi(\beta))$ , for  $\alpha < \beta \leq \omega_1$ ,
- (iii)  $\varphi(\omega_1) = x$ .

- In our case, there is  $\varphi: [0, \omega_1] \rightarrow \mathcal{K}_\varrho$  with

- (i)  $|\varphi(\alpha)| \leq \omega$ , for  $\alpha < \omega_1$ ,
- (ii)  $\varphi(\alpha) \subseteq \varphi(\beta)$ , for  $\alpha < \beta \leq \omega_1$ ,
- (iii)  $\varphi(\omega_1) = A, \implies \bigcup_{\alpha < \omega_1} \varphi(\alpha) = A$ .

- Hence, the sets  $\varphi(\alpha)$  are infinite, for  $\alpha$  large.

- By the previous case, such  $\varphi(\alpha)$  are initial intervals. ✓ ■



- ① Introduction and main results
- ②  $\varrho$ -functions and Theorem B
- ③ Proof of Theorem A**
- ④ Semi-Eberlein compacta (*very* briefly)



# Working in the space $C(\mathcal{K}_\rho)$

By Th B, pick  $\mathcal{F}_\rho \subseteq [\omega_1]^{<\omega}$  such that  $\mathcal{K}_\rho := \overline{\mathcal{F}_\rho}$  satisfies:

- (i)  $\{\alpha\} \in \mathcal{K}_\rho$  for every  $\alpha < \omega_1$ ,
- (ii)  $[0, \alpha) \in \mathcal{K}_\rho$  for every  $\alpha \leq \omega_1$ ,
- (iii) ... (*Sth we don't need anymore*)
- (iv)  $\mathcal{K}_\rho$  is scattered.

We define a biorthogonal system  $\{f_\gamma; \mu_\gamma\}_{\gamma < \omega_1}$  in  $C(\mathcal{K}_\rho)$ :

$$f_\gamma \in C(\mathcal{K}_\rho) \qquad f_\gamma(A) = \begin{cases} 1 & \gamma \in A \\ 0 & \gamma \notin A \end{cases} \quad (A \in \mathcal{K}_\rho)$$

$$\mu_\gamma := \delta_{\{\gamma\}} \in \mathcal{M}(\mathcal{K}_\rho) \qquad \mu_\gamma(S) = \begin{cases} 1 & \{\gamma\} \in S \\ 0 & \{\gamma\} \notin S \end{cases} \quad (S \subseteq \mathcal{K}_\rho).$$

- $\langle \mu_\alpha, f_\gamma \rangle = f_\gamma(\{\alpha\}) = \delta_{\alpha, \gamma}$ , so it is biorthogonal.



The space that we are looking for is

$$\mathcal{X}_Q := \overline{\text{span}}\{f_\gamma\}_{\gamma < \omega_1} \subseteq C(\mathcal{K}_Q).$$

What do we know already?

- $\mathcal{X}_Q$  is Asplund (as  $\mathcal{K}_Q$  is scattered);
- $\{f_\gamma; \mu_\gamma \upharpoonright \mathcal{X}_Q\}_{\gamma < \omega_1}$  is a biorthogonal system in  $\mathcal{X}_Q$ .

What do we still need?

- $\mathcal{X}_Q$  is not WLD;
- $\text{span}\{\mu_\gamma \upharpoonright \mathcal{X}_Q\}_{\gamma < \omega_1}$  is a 1-norming subspace for  $\mathcal{X}_Q$ ;
- In particular,  $\text{span}\{\mu_\gamma \upharpoonright \mathcal{X}_Q\}_{\gamma < \omega_1}$  is  $w^*$  dense, so  $\{f_\gamma; \mu_\gamma \upharpoonright \mathcal{X}_Q\}_{\gamma < \omega_1}$  is an M-basis for  $\mathcal{X}_Q$ .





- We shall show that  $[0, \omega_1]$  embeds in  $(B_{\mathcal{X}_\varrho^*}, w^*)$ .
- Define  $\iota: [0, \omega_1] \rightarrow \mathcal{K}_\varrho$  by  $\alpha \mapsto [0, \alpha)$  (recall that  $[0, \alpha) \in \mathcal{K}_\varrho$ ).

$$\begin{array}{ccccc}
 [0, \omega_1] & \xhookrightarrow{\iota} & \mathcal{K}_\varrho & \xhookrightarrow{\delta} & (B_{\mathcal{M}(\mathcal{K}_\varrho)}, w^*) \\
 & & \searrow \text{---} e \text{---} & & \downarrow q \\
 & & & & (B_{\mathcal{X}_\varrho^*}, w^*)
 \end{array}$$

- **Claim.**  $e$  is injective.
  - Note that  $e(A) := \delta_A \upharpoonright_{\mathcal{X}_\varrho}$ ;
  - Let  $A \neq B \in \mathcal{K}_\varrho$  and pick  $\gamma \in A \setminus B$ ;
  - $\langle \delta_A \upharpoonright_{\mathcal{X}_\varrho}, f_\gamma \rangle = \langle \delta_A, f_\gamma \rangle = f_\gamma(A) = 1$ ;
  - $\langle \delta_B \upharpoonright_{\mathcal{X}_\varrho}, f_\gamma \rangle = f_\gamma(B) = 0$ .





$\text{span}\{\mu_\gamma \upharpoonright \mathcal{X}_\varrho\}_{\gamma < \omega_1}$  is 1-norming

**Claim.** Let  $A \in \mathcal{F}_\varrho$ . Then

$$\delta_A \upharpoonright \mathcal{X}_\varrho = \sum_{\alpha \in A} \delta_{\{\alpha\}} \upharpoonright \mathcal{X}_\varrho.$$

*Proof.* Just check that  $\langle \delta_A, f_\gamma \rangle = \langle \sum_{\alpha \in A} \delta_{\{\alpha\}}, f_\gamma \rangle$ . ✓

In particular,

$$(\dagger) \quad \{\delta_A \upharpoonright \mathcal{X}_\varrho : A \in \mathcal{F}_\varrho\} \subseteq \text{span}\{\mu_\gamma \upharpoonright \mathcal{X}_\varrho\}_{\gamma < \omega_1}.$$

Finally, for every  $f \in \mathcal{X}_\varrho$  we have

$$\|f\| = \max_{A \in \mathcal{K}_\varrho} |f(A)| = \sup_{A \in \mathcal{F}_\varrho} |f(A)| = \sup_{A \in \mathcal{F}_\varrho} |\langle \delta_A, f \rangle|$$

$$\stackrel{(\dagger)}{\leq} \sup \left\{ |\langle \mu, f \rangle| : \mu \in \text{span}\{\mu_\gamma \upharpoonright \mathcal{X}_\varrho\}_{\gamma < \omega_1}, \|\mu\| \leq 1 \right\}.$$



- ① Introduction and main results
- ②  $\varrho$ -functions and Theorem B
- ③ Proof of Theorem A
- ④ Semi-Eberlein compacta (*very* briefly)




## Definition (Kubiś and Leiderman, 2004)


A compact space is **semi-Eberlein** if it is homeomorphic to a compact  $\mathcal{K} \subseteq [0, 1]^\Gamma$  such that  $c_0(\Gamma) \cap \mathcal{K}$  is dense in  $\mathcal{K}$ .

Recall that  $\Sigma(\Gamma) := \{x \in [0, 1]^\Gamma : \text{supp}(x) \text{ is countable}\}$ .

A compact space is ...



if it is homeomorphic to  $\mathcal{K} \subseteq [0, 1]^\Gamma$  such that ...





## Theorem (Kubiś and Leiderman, 2004)

No semi-Eberlein compact space has a P-point.

- Used to show that there is  $\mathcal{K}$  Corson, not semi-Eberlein.
- A point  $p \in \mathcal{K}$  is a **P-point** if it is not isolated and for every choice of  $(U_j)_{j < \omega}$  nghds of  $p$ ,  $\cap U_j$  is a nghd of  $p$ .

## Question (Kubiś and Leiderman, 2004)

Can a semi-Eberlein compact space have weak P-points?

- A point  $p \in \mathcal{K}$  is a **weak P-point** if it is not isolated and no sequence in  $\mathcal{K} \setminus \{p\}$  converges to  $p$ .
- The compact space  $\mathcal{K}_\mathcal{Q}$  in Theorem B is semi-Eberlein and it has a weak P-point.



### Theorem A

There exists an Asplund space  $X$  with a 1-norming M-basis such that  $X$  is not WLD.

### Theorem B

There exists a family  $\mathcal{F}_\rho \subseteq [\omega_1]^{<\omega}$  of finite subsets of  $\omega_1$  such that  $\mathcal{K}_\rho := \overline{\mathcal{F}_\rho}$  has the following properties:

- (i)  $\{\alpha\} \in \mathcal{K}_\rho$  for every  $\alpha < \omega_1$ ,
- (ii)  $[0, \alpha) \in \mathcal{K}_\rho$  for every  $\alpha \leq \omega_1$ ,
- (iii) if  $A \in \mathcal{K}_\rho$  is an infinite set, then  $A = [0, \alpha)$  for some  $\alpha \leq \omega_1$ ,
- (iv)  $\mathcal{K}_\rho$  is scattered.

**Thank you for your attention!**