

Lipschitz functions and polynomials on ℓ_2

Tommaso Russo

tommaso.russo.math@gmail.com

P. Hájek and T. Russo

Projecting Lipschitz functions onto spaces of polynomials

Mediterr. J. Math. **19** (2022), 190.

Æsy to define, Hard to analyse:

First conference on Lipschitz free spaces

Besançon, September 19–22, 2023

Are spaces of Lipschitz functions different?



Problem (Candido–Cúth–Doucha)

Are there two separable, infinite-dimensional Banach spaces \mathcal{X} and \mathcal{Y} such that $\text{Lip}_0(\mathcal{X})$ and $\text{Lip}_0(\mathcal{Y})$ are not isomorphic?

- ▶ **Candido–Kaufmann ('21).** $w^*\text{-dens}(\text{Lip}_0(\mathcal{X})^*) = \text{dens}(\mathcal{X})$.
- ▶ $\text{Lip}_0(\mathbb{R}) \simeq \ell_\infty$.
- ▶ **Kisliakov ('75), Cúth–Doucha–Wojtaszczyk ('16).**
If $\dim(\mathcal{X}) \geq 2$, there is no onto map from a $\mathcal{C}(\mathcal{K})$ space to $\text{Lip}_0(\mathcal{X})$.
- ▶ Is $\text{Lip}_0(\mathbb{R}^k) \simeq \text{Lip}_0(\mathbb{R}^n)$, for $k, n \geq 2$ distinct?
- ▶ Is there an infinite-dimensional \mathcal{X} with $\text{Lip}_0(\mathcal{X}) \simeq \text{Lip}_0(\mathbb{R}^2)$?
- ▶ **Candido–Cúth–Doucha ('19).** If \mathcal{X} is a separable, inf.-dim. \mathcal{L}_p -space, then $\text{Lip}_0(\mathcal{X}) \simeq \text{Lip}_0(\ell_p)$ when $p < \infty$ and $\text{Lip}_0(\mathcal{X}) \simeq \text{Lip}_0(c_0)$ when $p = \infty$.
 - ▶ **Dutrieux–Ferenczi ('05).** The case $\mathcal{X} = \mathcal{C}(\mathcal{K})$.
- ▶ **Candido–Cúth–Doucha ('19).** Is $\text{Lip}_0(L_p) \simeq \text{Lip}_0(L_q)$ for $p \neq q$?



- ▶ A Banach space \mathcal{X} has the **approximation property (AP)** if for every compact set $\mathcal{K} \subseteq \mathcal{X}$ and $\varepsilon > 0$ there exists a finite-rank operator $T: \mathcal{X} \rightarrow \mathcal{X}$ such that $\|Tx - x\| < \varepsilon$ ($x \in \mathcal{K}$).
- ▶ \mathcal{X} has the **λ -BAP** if additionally $\|T\| \leq \lambda$. **MAP** \equiv **1-BAP**.
- ▶ AP and BAP pass to complemented subspaces and from \mathcal{X}^* to \mathcal{X} .
- ▶ **Johnson ('75)**. If $\text{Lip}_0(\mathcal{X})$ has AP/BAP, then \mathcal{X} has it too.
 - ▶ **Lindenstrauss ('64)** \mathcal{X}^* is 1-complemented in $\text{Lip}_0(\mathcal{X})$.
- ▶ However, the converse fails.
 - ▶ There exists \mathcal{X} with MAP such that \mathcal{X}^* fails AP.
- ▶ So, just take \mathcal{X} infinite-dimensional such that $\text{Lip}_0(\mathcal{X})$ has AP.
- ▶ Well, "just"...
- ▶ **Godefroy**. Does $\text{Lip}_0(\ell_2)$ have the AP?
- ▶ Actually, does $\text{Lip}_0(\mathbb{R}^n)$ have the AP/BAP, for $n \geq 2$?
- ▶ **Godefroy–Kalton ('03)**. $\mathcal{F}(\mathcal{X})$ has λ -BAP if and only if \mathcal{X} has it.



Problem (Godefroy)

Does $\text{Lip}_0(\ell_2)$ have the AP?

- ▶ Let's try to disprove it, judiciously.
- ▶ Candidates complemented subspaces of $\text{Lip}_0(\ell_2)$ that fail the AP?
 - ▶ **Szankowski ('81)**. $\mathcal{L}(\ell_2)$ fails the AP.
 - ▶ **Floret ('97), Dineed–Mujica ('15)**. $\mathcal{P}({}^2\ell_2)$ fails the AP.
 - ▶ In fact, $\mathcal{P}({}^2\ell_2) \simeq \mathcal{B}(\ell_2)$.
- ▶ $\mathcal{P}({}^2\mathcal{X})$ is the space of bounded 2-homogeneous polynomials on \mathcal{X} .
 - ▶ $P \in \mathcal{P}({}^2\mathcal{X})$ if there is a bounded bilinear map $M: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ such that $P(x) = M(x, x)$;
 - ▶ There is a unique symmetric such bilinear map;
 - ▶ $\|P\|_{\mathcal{P}} = \sup_{x \in B_{\mathcal{X}}} |P(x)|$.
- ▶ But polynomials are **not** Lipschitz functions!



- ▶ However, they are Lipschitz on the unit ball.
 - ▶ Therefore, $\mathcal{P}(^2\mathcal{X})$ is a natural subspace of $\text{Lip}_0(B_{\mathcal{X}})$.
 - ▶ Moreover, $\|\cdot\|_{\mathcal{P}}$ is equivalent to $\|\cdot\|_{\text{Lip}}$.
- ▶ Consequently, $\mathcal{P}(^2\mathcal{X})$ is naturally isomorphic to a subspace of $\text{Lip}_0(B_{\mathcal{X}})$, via the restriction map

$$P \mapsto P|_{B_{\mathcal{X}}}.$$

(Another) Problem

Is $\mathcal{P}(^2\ell_2) \subseteq \text{Lip}_0(B_{\ell_2})$ a complemented subspace?

- ▶ **Kaufmann ('15).** $\text{Lip}_0(\mathcal{X}) \simeq \text{Lip}_0(B_{\mathcal{X}})$.

Summary:

If $\mathcal{P}(^2\ell_2) \subseteq \text{Lip}_0(B_{\ell_2})$ is complemented, then $\text{Lip}_0(\ell_2)$ fails to have the AP.



Problem (*Repetita iuvant*)

Is $\mathcal{P}({}^2\ell_2) \subseteq \text{Lip}_0(B_{\ell_2})$ a complemented subspace?

- ▶ **Lindenstrauss ('64).** $\mathcal{X}^* = \mathcal{P}({}^1\mathcal{X})$ is 1-complemented in $\text{Lip}_0(\mathcal{X})$.
- ▶ No wonder, $\mathcal{P}({}^d\mathcal{X}) \equiv \text{bdd}, d\text{-homogeneous poly on } \mathcal{X}$.
- ▶ Is there a version of Lindenstrauss' result for polynomials?
- ▶ **Pełczyński ('57).** Every d -homogeneous polynomial from ℓ_p to ℓ_q is compact if $dq < p$.
- ▶ Polynomial Schur and Dunford-Pettis properties.
- ▶ Norm-attaining polynomials.
- ▶ Hahn-Banach extensions of polynomials.
- ▶ Can you, **please, state** something?



Theorem (Hájek, R., '22)

$\mathcal{P}({}^2\ell_2) \subseteq \text{Lip}_0(B_{\ell_2})$ is not complemented.

- ▶ We still don't know if $\text{Lip}_0(\ell_2)$ has the AP.
 - ▶ Maybe, that's an indication that it does.
 - ▶ At least, we **know** that this is not the correct approach.

Theorem (Hájek, R., '22)

If \mathcal{X} contains uniformly complemented $(\ell_2^n)_{n=1}^\infty$, then $\mathcal{P}({}^2\mathcal{X})$ is not complemented in $\text{Lip}_0(B_{\mathcal{X}})$.

- ▶ In particular, if \mathcal{X} has non-trivial type...
- ▶ Moreover, for every $d \geq 2$, $\mathcal{P}({}^d\mathcal{X})$ and $\mathcal{P}_0^d(\mathcal{X})$ are not complemented in $\text{Lip}_0(B_{\mathcal{X}})$.
- ▶ $\mathcal{P}^d(\mathcal{X})$ is the space of polynomials of degree at most d on \mathcal{X} .



- ▶ **Problem.** Is $\mathcal{P}({}^2c_0)$ complemented in $\text{Lip}_0(B_{c_0})$?
 - ▶ **Alencar ('75).** $\mathcal{P}({}^2c_0)$ has a Schauder basis (so BAP).
- ▶ **Aron–Schottenloher ('76).** $\mathcal{P}({}^d\ell_1) \simeq \ell_\infty$.
 - ▶ $\mathcal{L}({}^d\ell_1) \simeq \ell_\infty$ (a computation).
 - ▶ $\mathcal{P}({}^d\ell_1)$ is complemented in $\mathcal{L}({}^d\ell_1)$ (Polarisation formula).
 - ▶ **Lindenstrauss ('67).** ℓ_∞ is prime.
- ▶ **Arias–Farmer ('96).** If \mathcal{X} is a separable \mathcal{L}_1 -space, $\mathcal{P}({}^d\mathcal{X}) \simeq \ell_\infty$.

(Main) Proposition

Let E be \mathbb{R}^n with Euclidean norm. If Q is any projection from $\text{Lip}_0(B_E)$ onto $\mathcal{P}({}^2E)$, then

$$\|Q\| \geq c \cdot n^{1/5}.$$

Thank you for your attention!