



Total Synthesis of Hyperforin


Ting, C. P.; Maimone, T. *J. Am. Chem. Soc.* **2015**, *137*, 10516







1 – 5



6 – 10



- 1) **1**, CuBr·DMS, LiCl, TMSCl
- 2) MeLi·LiI *then* **2**, HMPA *then* TsOH, Δ
- 3) LDA, prenyl bromide
- 4) LTMP, **3**
- 5) TMSCH₂N₂



Lithium tetramethyl piperidine (LTMP)

- 6) PhI(OAc)₂, KOH, MeOH
- 7) LTMP, TsCl
- 8) LTMP, *i*-PrCOCN,
- 9) *i*-PrMgBr·LiCl, LDA, Li(2-Th)CuCN, *then* prenyl bromide
- 10) LiCl, DMSO, Δ

What are the two major types of cuprates?

Gilman - R₂CuLiX or R₂CuMgX

Cyanocuprates - RCu(CN)Li – Less reactive than Gilman cuprates

Lipshutz - R₂Cu(CN)Li₂ – Higher order cuprates – More reactive – CN acts as dummy ligand

Hint: in step 2 there is also an isomerization

Propose a mechanism for step 4

What is the pK_a of LTMP?

pK_a = 35 for reference LDA is 36

Only A is taken forward. B can be recycled, how could this be done (2 steps)

NaOH, then TMSCH₂N₂ again.


Propose a mechanism for step 6

Suggest a reason that LDA was added in step 9.

Citation 22 in the main text suggests *i*-PrCl formed as a byproduct which consumes the magnesiate in an E₂ elimination reaction.

LDA is used to suppress this.

Mechanism for step 4:



Mechanism for step 6:

