



## Asymmetric Total Synthesis of (+)-Mannolide C

Qiaoqiao Ao, Hai-Jun Zhang, Jinbin Zheng, Xiaoming Chen, Hongbin Zhai  
*Angew. Chem. Int. Ed.* **2021**, *60*, 21267-21271.



1. **1**, 170 °C
2. MeMgBr, CuI, -20 °C then Ac<sub>2</sub>O, NaOAc, 60 °C
3. LiAlH(O*t*-Bu)<sub>3</sub>, -78 °C to -20 °C
4. MsCl, Et<sub>3</sub>N, DMAP, 0 °C
5. NaBH<sub>4</sub>, CeCl<sub>3</sub>, 0 °C
6. vinyl acetate, Lipozyme TL IM, 40 °C, 4d
7. DMP, NaHCO<sub>3</sub>, 0 °C




8. **2**, CuBr•DMS, HMPA, -78 °C then **A**, TMSCl, -20 °C
9. **3**, KHMDS, 0 °C then product of step 8, 60 °C then 2 M HCl, 40 °C
10. TMSOTf, (TMSOCH<sub>2</sub>)<sub>2</sub> 0 °C then *m*-CPBA, NaHCO<sub>3</sub> 0 °C
11. O<sub>3</sub>, -78 °C then DMS then **4**, KHMDS, -40 °C
12. Zn, AcOH, NaOAc, NaI, r.t. then conc. HCl, r.t.
13. **5**, CeCl<sub>3</sub>, -78 °C then DMP, 0 °C

Step 1: Name the reaction

Hint: Steps 5-6 are for resolution


Step 5: Name the reaction

Step 6: How does the resolution with Lipozyme TL IM work?




Hint: Two equivalents of **4** are used in step 11




**B**



14-22



14. **6**, MW, toluene, 150 °C
15. DBU, toluene, reflux
16. AlMe<sub>3</sub>, Ni(acac)<sub>2</sub>, LiBr, -5 °C
17. NaBH<sub>4</sub>, r.t. *then* K<sub>2</sub>CO<sub>3</sub>
18. Mn(OAc)<sub>3</sub>·2H<sub>2</sub>O, TBHP, 3Å MS, 55 °C
19. KHMDS, MeI -78 °C to 50 °C
20. TsNHNH<sub>2</sub>, toluene, 60 °C *then* silica gel, **7**, 0 °C *then* NaOAc, reflux
21. OsO<sub>4</sub>, NMO, TsOH, r.t.
22. **8**, toluene, reflux



Step 14: Please name the reagent

Step 22: Please name the reagent