



Total Synthesis of (+)-Fendleridine (Aspidioalbidine) and (+)-1-Acetylaspidoalbidine

E.L. Campbell, M. Zuhl, C.M. Liu, D.L. Boger

J. Am. Chem. Soc. 2010, 132, 9, 3009–3012




1-5



6-9

- 1) phthalic anhydride
- 2) NaH, BnBr
- 3) $N_2H_4 \cdot H_2O$
- 4) CDI
- 5) **1**, AcOH
- 6) TsCl, NEt₃



- 7) EDCI, DMAP, **2**
- 8) *o*-Cl₂C₆H₄, 180 °C
- 9) NH₃, MeOH then TFAA, pyridine
- 10) HF-pyridine

Name of starting material?


Tryptamine


6) Hint: an 1,3,4-oxadiazole is formed

8) Hint: Tandem [4+2] / [3+2] cycloaddition


9) Hint: loss of signals at $\delta = 171.2$ ppm (¹³C-NMR) and $\delta = 3.66$ ppm (s, ¹H-NMR)

10) two reactions occur (a cyanohydrine is formed)





10-15



(+)-Fendleridine

- 11) Na-selectride
- 12) NaH, CS₂ then MeI
- 13) AIBN, Bu₃SnH
- 14) Lawesson's reagent
- 15) Raney-Ni
- 16) Na, NH₃, t-BuOH

11) Hint: C₂₆H₂₈N₂O₃

12 & 13) Name of reaction?
Barton McCombie deoxygenation

14) Structure of Lawesson's reagent?

