


## Enantioselective Total Synthesis of (+)-Salvileucalin B

Sergiy Levin, Roger R. Nani, and Sarah E. Reisman.


*J. Am. Chem. Soc.* **2011**, 133, 774-776.



1 - 4



**A**





5 - 6



**B**


- 1) Me<sub>2</sub>Zn, **1**, PhMe, 70 °C then 3-furaldehyde, 0 to 22 °C
- 2) NaH, **2**, DMF
- 3) 1M HCl, MeOH
- 4) MsCl, NEt<sub>3</sub>, THF then LiBr



- 5) LiHMDS, LiCl, THF, -78 to 22 °C then **A**, -78 °C
- 6) TBAF, CH<sub>2</sub>Cl<sub>2</sub>

1) Name of chiral ligand **1**?

5) Name of the auxiliary on the starting material? Who developed this chemistry?




**(+)-Salvileucalin B**

**B**




7-13



**C**




14-18



**(+)-Salvileucalin B**

- 7)  $\text{RuCp}^*(\text{cod})\text{Cl}$  (8 mol%),  $\text{CH}_2\text{Cl}_2$ , 45 °C
- 8)  $n\text{-Bu}_4\text{NOH}$ ,  $t\text{-BuOH}/\text{H}_2\text{O}$ , 90 °C
- 9)  $(\text{COCl})_2$ , DMF (cat.) then  $\text{CH}_2\text{N}_2$ , THF
- 10)  $\text{AgTFA}$ , MeOH,  $\text{NEt}_3$ , THF, -30 to 22 °C
- 11)  $\text{NaCH}_2\text{CN}$ , THF, -78 to 22 °C
- 12)  $(\text{imid})\text{SO}_2\text{N}_3$ , pyridine
- 13)  $\text{Cu}(\text{hfacac})_2$  (10 mol%),  $\text{CH}_2\text{Cl}_2$ , 120 °C, microwave

- 14)  $\text{NaHMDS}$ , -78 °C then  $\text{Ti}_2\text{NPh}$
- 15)  $\text{DIBAL}$ ,  $\text{CH}_2\text{Cl}_2$ , -40 °C then  $\text{AcOH}$  aq.
- 16)  $\text{DIBAL}$ ,  $\text{CH}_2\text{Cl}_2$ , -40 °C then  $\text{AcOH}$  aq.
- 17)  $\text{Pd}_2(\text{dba})_3$  (5 mol%),  $\text{dppf}$  (10 mol%), CO, DIPEA, THF
- 18) **3**,  $\text{CH}_2\text{Cl}_2$ , -35 °C



7) *Hint*: 3 rings are formed.

10) Name of the reaction?

15) Name of the reaction?